

Effect of Date and Quantity of Potassium Fertilizer Application on Growth, Yield and Oil Quality of Sunflower *Helianthus annuus L. var. flame*

Faisal, M. Al –Taher¹, Yahya, K. Jellab¹, Saad, A. M. Al-badry²

¹ Field crop Department, College of Agriculture/ University of Al- Muthanna

² Field crop Department, College of Agriculture/ University of Thi-qar

Abstract: A field experiment was carried out in Thi-qar province – Batha city with two Seasons, 1st experiment was during Fall season (2011) and 2nd experiment was during Spring Season (2012) using R.C.B.Desgn. with three replications. The main objective of experiment was to find the effect of dates and levels of Potassium on oil quality and yield of Sunflower. Six levels of K were used (K_0 = Control, $K_1= 120 \text{ kgK.ha}^{-1}$ (Soil), $K_2= 6000 \text{ mgk.L}^{-1}$ foliar, $K_3=30\text{kgk.ha}^{-1}$ soil+ 3000mgk.L^{-1} foliar, $K_4=30\text{kgk.ha}^{-1}$ soil+ 6000mgk.L^{-1} foliar, $K_5=30\text{kgk.ha}^{-1}$ soil + 9000 mgk.L^{-1} foliar) and three dates ($T_1= 8$ leaves, $T_2= 14$ leaves, $T_3= 20$ leaves). The results showed that increasing of Potassium levels significantly caused increasing in (seed number, 1000 seed weight and yield plant) Fall and Spring seasons. The highest seed yield was 28,7% and it obtained by third date of 2nd season. Also the percent of oil and oil yield of 1st and 2nd season were increased respectively. The interaction had significant effect on yield traits (seed number and seed 1000 weight) of 2nd season. There was no effect for fertilization and dates in oleic acid for both seasons, while the linoleic acid was influenced by 1st date of 1st and 2nd season respectively

Key words- Sunflower seed yield. Potassium application. Application date. Application quantity. Oil quality.

I. Introduction

Sunflower (*Helianthus annuus L.*) is an important oil seed crop in the world and it belongs to the family Asteraceae. It increased in last years as a result of the decreasing in the quantity of produced oils because of using it in many fields of life. In addition, the seeds contain high percentage of protein ranging from 20-30%, and it used in poultry feed as well [1], Also the seeds contain high percent of oil reach to 50% [2]. Sunflower oil characterizes with low saturated fatty acids that have essential role in arteries intransigence and high content of E vitamin [3]. The productivity of any crop is mainly based on several factors in which the agricultural practices come in at first, including the quantity and date of fertilizers application which consider important matters that can enable the crop to exploitation of different environmental factors (light, water, soil, etc.) with best way and increase the quantity and quality of yield per area, so many researchers have turned to use various means to increase the production by means of adding the micro and macro elements, particularly the potassium element, which plays an important role in increase and improve the quality of yield [4]. The researcher of Iraqi cropping systems notice careless of using potassium fertilizers for long time due to popular misconception that Iraqi soils are rich with potassium element, however fact this element exposes to fixation process in the primary and secondary clay minerals which change it from soluble or exchangeable form to slowly available in the soil solution [5].

Practically plant can't grow and develop normally with out of potassium element, so it can contributes in many important physiological processes in plant for instances, regulation of osmotic pressure of plant cells and it can controls the process of opening and closing of stomates. It also has a relationship to synthesis of protein and fats, synthesis of chlorophyll, stimulation of adenosine triphosphate (ATP) formation and photosynthesis, anabolism and metabolism of carbohydrates, translocation of photosynthetic. In addition potassium used to increase of lipids synthesis in oil crops. It also can contribute in action of approximately 60 enzymes [7].

So as showed above and on the basis of benefits of this crop as well as the positive effect of fertilization to increasing the yield per area with improving its quality, besides the scarcity of researches especially about potassium fertilization in Iraq, so this study was carried out in order to determination of best concentration of fertilizer for obtaining best productivity and quality per area and best time for potassium application.

II. Materials and methods

The field experiment was carried out during two seasons fall (2011) and spring (2012) at Batha district to study the effect of date and quantity of potassium fertilizer application on growth, yield and oil quality of Sunflower. The experiment was laid out in a Random Complete Block Design (R.C.B.D) with three replications.

Soil samples were taken randomly from different locations for the entire experiment a study area in both seasons within layer (0-30) cm. It had dried, crushed and sieved with sieve has mesh with (2mm) diameter then mixed together. A complex sample was taken for each season. Physical and Chemical tests were done pre-cultivation including electrical conductivity (EC) and soil pH) according to [8], soil texture and available potassium as it revealed in Table (1).

Table (1) Mechanical and Chemical Analysis during Fall and Spring seasons 2011 - 2012

Soil Analysis	Fall Season		Spring Season	
pH	8.3		7.9	
Ec ms/cm ⁻¹	3.6		3.9	
Soil texture	Clay		Clay	
Available K (PPm)	Pre – Cultivation	Post - Cultivation	Pre – Cultivation	Post - Cultivation
	165	155	167	156

The researched factors in the experiment were six levels of potassium and three dates of application:

A. Potassium levels:

- K₀: control.
- K₁: 120 kg K. ha⁻¹soil.
- K₂: 6000 mg K .L⁻¹ Foliar spray.
- K₃: 30 kg K. ha⁻¹ soil + 3000 mg K .L⁻¹Foliar spray.
- K₄: 30 kg K .ha⁻¹ soil + 6000 mg K .L⁻¹ Foliar spray.
- K₅: 30 kg K .ha⁻¹ soil + 9000 mg K .L⁻¹ Foliar spray.

B. Dates of application:

- T1= 8 true leaves.
- T2= 14 true leaves.
- T3= 20 true leaves.

The researched attributes were (number of seeds, 1000 seeds weight, seed yield of plant, Total seed yield, oil percent and Oil yield). All data were analyzed statically using (R.C.B.D.) and the means compared with (R.L.S.D.) at (P≤ 0.05) using (Gen stat v.10.3) software.

III. Results and discussion

1. Fall season

1.1 Seed Number (seed.head⁻¹)

The data presented in table (2) indicated that potassium application significantly affected the seed number. head⁻¹. Maximum seed number (861) seed.head⁻¹ was observed on plants that treated with (K₁) treatment, whereas minimum value (589) seed.head⁻¹ was observed on plants that treated with control (K₀) treatment. K₃ and K₅ treatments had non-significant effect on this trait between them. Our results is in agreement with [9] May be due to the role of fertilization, environmental conditions and variety during formation stage and completion of flowering buds, so the seed number was increased significantly.

Also with respect to fertilization dates had significant effect on seed number. The Second date (T₂) gave maximum seed number (867) seed.head⁻¹, whereas minimum value (636) seed.head⁻¹ was recorded at First date (T₁) due to potassium and other elements are existing in sufficient quantities during seeds formation. The result is in agreement with [1]. In addition, the interaction effect between quantity and date of fertilization table (3) showed significant differences on seed number, so (K₅T₂) treatment gave maximum seed number (1006) seed.head⁻¹, whereas minimum value (392) seed.head⁻¹ was recorded with (K₀T₂) treatment .

1.2 1000 Seed weight (gm)

The data presented in table (2) indicated that potassium application significantly affected the 1000 seed weight. Maximum 1000 Seed weight (77) gm was observed on plants that treated with (K₄) treatment, whereas minimum value (48.5) gm was observed on plants that treated with (K₃) treatment. K₁, K₂ and K₅ treatments had non-significant effect on this trait between them. The result is in agreement with [3]. This increasing may be due to the efficiency of plants to translocation of photosynthates towards seeds resulting in more filled seeds (source/ sink efficiency). The interaction effect between quantity and date of fertilization table (3) showed significant differences on 1000 seed weight, so (K₀T₁) treatment gave maximum 1000 Seed weight (124.3) gm, whereas minimum value (44.2) gm was recorded with (K₀T₃) treatment that on a par with (K₁T₁) treatment.

1.3 plant yield (gm)

The data presented in table (2) indicated that potassium application significantly affected the plant yield. Maximum plant yield (67.9) gm was observed on plants that treated with (**K₄**) treatment, whereas minimum value (39.6) gm was observed on plants that treated with (**K₃**) treatment. **K₀**, **K₁**, **K₂**, and **K₅** treatments had non-significant effect on this trait between them. This result is in agreement with [9, 13 and 15] . It is possibly due to the role of fertilization for increasing photosynthesis and translocation of photosynthetic from source to sink. With respect to fertilization dates were affected by plant yield of seeds. The second date (**T₂**) gave maximum yield (59.5) gm.plant⁻¹, whereas minimum value (39) gm.plant⁻¹ was recorded at first date (**T₁**). The result is in agreement with [13] due to existence of potassium element in sufficient quantities during growth stages.

1.4 Seed yield (ton.ha⁻¹)

The data presented in table (2) indicated that potassium application significantly affected the seed yield. Maximum seed yield (4.36) ton.ha⁻¹ was observed on plants that treated with (**K₁**) treatment which was superior to all the other treatments, whereas minimum value (2.30) ton.ha⁻¹ was observed on plants that treated with control (**K₅**) treatment. The result is in agreement with [9 and 10]. It is possibly due to the role of fertilization for increasing plant yield resulted in increase of seeds yield. Seed yield was affected significantly by fertilization dates. Third date gave maximum seed yield (3.65) ton.ha⁻¹ which was superior to the second date (2.60) ton.ha⁻¹ This result is in agreement with [9] due to fertilization date was on time and with optimal level during yield formation.

1.5 Oil content (%)

The data presented in table (2) indicated that fertilization date significantly affected the oil content. Third date (**T₃**) gave maximum oil content (51.1)% which was superior to all the other dates. The result is in agreement with [9, 13] the increasing of potassium application at third date with light and temperature as well as role of potassium in structure of fatty acids may be caused more accumulation of oil towards seeds..

1.6 Oil yield ton.ha⁻¹

The data presented in table (2) indicated that fertilization date significantly affected the oil yield. Third date (**T₃**) gave maximum oil yield (1.907) ton.ha⁻¹ which was superior to all the other dates, whereas minimum value (1.058) ton.ha⁻¹ was recorded at second date (**T₂**). The result is in agreement with [9] due to application date of potassium was sufficient and on time during that stage to increase oil yield.

1.7 Oleic acid (%)

The data presented in table (4) indicated that quantity and date of potassium application, and interaction between them had non-significant effect on oleic acid. The result is contradictory to [12].

1.8 Linoleic acid (%)

The data presented in table (4) indicated that quantity of potassium application had significant effect on linoleic acid. The treatment (**K₁**) gave maximum percentage of linoleic acid (0.62) % which was on a par with (**K₃** and **K₅**) treatments, whereas minimum value (0.36) % was recorded with (**K₀**) treatment. This result is in agreement with [10, 11] and contradictory to [12].

1.9 Palmitic acid (%)

The data presented in table (4) indicated that dates of potassium application had significant effect on palmitic acid. First date (**T₁**) gave maximum percentage (7.96) % which was superior to all the other dates, whereas minimum value (4.62) % was recorded at third date (**T₃**). The result is in agreement with [12].

1.10 Stearic acid (%)

The data presented in table (4) indicated that quantity and date of potassium application, and interaction between them had non-significant effect on stearic acid. The result is contradictory to that reported by [12] who found that application of potassium reduced stearic acid percentage.

1.11 Myristic acid (%)

The data presented in table (4) indicated that quantity, date of potassium application, and interaction between them had non-significant effect on myristic acid.

2. Spring season

2.1 Seed Number (seed.head⁻¹)

The data presented in table (6) indicated that potassium application significantly affected the seed number.head⁻¹. Maximum seed number (861) seed.head⁻¹ was observed on plants that treated with (K₁) treatment which was superior to all the other dates, whereas minimum value (597) seed.head⁻¹ was observed on plants that treated with (K₄) treatment. This result is in agreement with [9, 14] may be due to the role of fertilization, environmental conditions and variety during formation stage and completion of flowering buds, so the seed number was increased significantly.

2.2 1000 Seed weight (gm)

The data presented in table (6) indicated that potassium application significantly affected the 1000 seed weight. Maximum 1000 seed weight (84.2) gm was observed on plants that treated with (K₃) treatment, whereas minimum value (31.4) gm was observed on plants that treated with (K₅) treatment. The result is in agreement with [3] This increasing may be due to the efficiency of plants to translocation of photosynthates towards seeds resulting in more filled seeds (source/ sink efficiency)..

2.3 plant yield (gm)

The data presented in table (6) indicated that potassium application significantly affected the plant yield. Maximum plant yield (55.9) gm was observed on plants that treated with (K₃) treatment, whereas minimum value (26.1) gm was observed on plants that treated with (K₅) treatment. The result is in agreement with [9,13 and 15] possibly due to the role of fertilization for increasing photosynthesis and translocation of photosynthetic from source to sink.

2.4 Seed yield (ton.ha⁻¹)

The data presented in table (6) indicated that potassium application significantly affected the seed yield. Maximum Seed yield (3.72) ton.ha⁻¹ was observed on plants that treated with (K₃) treatment which was superior to all the other treatments, whereas minimum value (1.74) ton.ha⁻¹ was observed on plants that treated with (K₅) treatment. The result is in agreement with [9 and 13] it is may be due to the role of fertilization for increasing plant yield resulted in increase of seeds yield.

2.5 Oil content (%)

The data presented in table (6) indicated that fertilization date significantly affected the oil content. Third date (T₃) gave maximum oil content (16.65)% which was superior to all the other dates whereas minimum value (13.19) % was observed with first date (T₁). The result is in agreement with [9, 13] the increasing of potassium application at third date with light and temperature as well as role of potassium in structure of fatty acids may be caused more accumulation of oil towards seeds..

2.6 Oil yield ton.ha⁻¹

The data presented in table (6) indicated that fertilization date significantly affected the oil yield. Third date (T₃) gave maximum oil yield (0.492) ton.ha⁻¹ which was superior to all the other dates, whereas minimum value (0.297) ton.ha⁻¹ was recorded at second date (T₂). The result is in agreement with [9] possibly due to of potassium application was sufficient and on time during that stage to increase oil yield.

2.7 Oleic acid (%)

The data presented in table (8) indicated that quantity and date of potassium application, and interaction between them had non-significant effect on oleic acid. The result is contradictory to [12].

2.8 Linoleic acid (%)

The data presented in table (8) indicated that quantity of potassium application had significant effect on linoleic acid. The treatment (K₁) gave maximum percentage of linoleic acid (0.62) % which was superior to all the other treatments, whereas minimum value (0.36) % was recorded with (K₀) treatment which was on a par with (K₂ and K₄) treatments, The result is in agreement with [10, 11].

Also, dates of potassium application had significant effect on linoleic acid. Second date (T₂) gave maximum percentage (0.56) % which was superior to all the other dates, whereas minimum value (0.37) % was recorded at first date (T₁). The result is in agreement with [12]. With respect the interaction between dates and quantities of potassium application had significant effect on linoleic acid table (9). The treatment (K₁T₂) gave maximum percentage of linoleic acid (0.70) % which was superior to all the other treatments, whereas minimum value (0.12) % was recorded with (K₀T₁) treatment.

2.9 Palmitic acid (%)

The data presented in table (8) indicated that dates of potassium application had significant effect on palmitic acid. First date (T₁) gave maximum percentage (8.90) % which was superior to all the other dates, whereas minimum value (3.95) % was recorded at third date (T₃). The result is in agreement with [12].

With respect the interaction between dates and quantities of potassium application had significant effect on palmitic acid table (9). The treatment (K₀T₁) gave maximum percentage of palmitic acid (9.65) % which was on a par with (K₄T₁), whereas minimum value (1.90) % was recorded with (K₂T₃) treatment.

2.10 Stearic acid (%)

The data presented in table (8) indicated that and dates of potassium application, had significant effect on stearic acid. Second date (T₂) gave maximum percentage (7.21) % which was superior to all the other dates, whereas minimum value (1.90) % was recorded at third date (T₃). This result is in agreement with [12].

2.11 Myristic acid (%)

The data presented in table (9) indicated that the interaction between dates and quantities of potassium application had significant effect on myristic acid. The treatment (K₀T₂, K₁T₁, K₂T₂, K₃T₁, K₄T₂ and K₅T₁) gave maximum percentage of myristic acid (0.07) %, whereas minimum value (0.02) % was recorded with (K₂T₃) treatment

Table (2) Effect of dates and quantity of potassium application on yield traits during fall season 2011

K treatments	No. seeds Seed.head ⁻¹	1000-seed weight (gm)	Plant yield (gm)	Total seed yield (ton.ha ⁻¹)	Oil content (%)	Oil yield (ton.ha ⁻¹)
K0	589	75.2	43.4	2.89	47.0	1.633
K1	861	51.1	46.4	3.09	43.2	1.569
K2	802	58.6	45.6	3.03	46.0	1.497
K3	774	48.5	39.6	2.63	42.8	1.888
K4	853	77.0	67.9	4.52	44.0	1.178
K5	751	58.5	46.3	3.08	40.8	0.941
L.S.D 0.05	80.58	17.87	15.87	1.057	N.S.	N.S.
Dates	636	64.6	39.0	2.60	40.7	1.387
T1						
T2	867	65.5	59.5	3.96	40.1	1.058
T3	811	54.4	46.1	3.07	51.1	1.907
L.S.D 0.05	56.9	N.S.	11.22	0.747	6.06	0.476

Table (3) Effect of interactions between Dates and Potassium Quantities on Yield traits during Fall season.

Traits	Dates	Potassium Quantities					
		K0	K1	K2	K3	K4	K5
Number of Seeds	D1	392	827	640	674	727	556
	D2	719	871	393	781	934	1006
	D3	657	884	872	866	900	689
	L.S.D (0.05)	139.4					
1000 Seeds Weight (gm)	D1	124.3	44.2	51.3	43.2	64.5	60.3
	D2	57.2	62.8	70.2	52.1	86.0	64.6
	D3	44.2	46.3	54.4	50.2	80.4	50.6
	L.S.D (0.05)	30.95					
Seed Yield of plant (gm)	D1	48.8	43.0	26.2	29.2	51.1	35.8
	D2	52.7	54.8	62.7	41.0	80.3	65.5
	D3	28.9	41.4	47.8	48.5	72.4	37.6
	L.S.D (0.05)	N.S.					
Total seed Yield (ton.ha ⁻¹)	D1	3.25	2.86	1.74	1.94	3.40	2.38
	D2	3.51	3.65	4.17	2.73	5.35	4.36
	D3	1.92	2.75	3.18	3.23	4.82	2.50
	L.S.D (0.05)	N.S.					
Oil percent %	D1	40.4	42.8	39.7	42.6	37.7	40.6
	D2	38.9	43.5	38.3	42.8	36.3	40.8
	D3	61.5	43.4	60.0	42.9	58.0	40.8
	L.S.D (0.05)	N.S.					
Oil yield	D1	1.090	1.669	1.264	1.336	0.715	1.249
	D2	1.228	1.344	1.000	1.230	0.773	0.774
	D3	2.580	1.695	2.225	2.098	2.045	0.799
	L.S.D (0.05)	N.S.					

Table (4) Effect of Dates and Quantity of Potassium Application on Fatty acids (%) during Fall Season 2011

K treatments	Fatty acids contents (%)				
	Oleic (%)	Linoleic (%)	Palmitic (%)	Stearic (%)	Myristic (%)
K0	0.026	0.36	6.68	5.00	0.04
K1	0.023	0.62	6.37	6.73	0.06
K2	0.018	0.45	5.83	4.83	0.05
K3	0.023	0.62	6.37	6.73	0.06
K4	0.023	0.43	5.87	5.13	0.05
K5	0.014	0.62	5.83	5.83	0.06
L.S.D 0.05	N.S.	0.203	N.S.	N.S.	N.S.
Dates					
T1	0.034	0.43	7.96	6.47	0.05
T2	0.021	0.59	5.89	6.17	0.06
T3	0.009	0.53	4.62	4.47	0.05
L.S.D 0.05	N.S.	N.S.	1.838	N.S.	N.S.

Table (5) Effect of interactions between Dates and Potassium Quantities on Fatty Acids content (%) during fall season 2011

Fatty acids	Dates	Potassium Quantities					
		K0	K1	K2	K3	K4	K5
Oleic (%)	D1	0.030	0.045	0.031	0.045	0.031	0.020
	D2	0.045	0.019	0.004	0.019	0.020	0.019
	D3	0.002	0.005	0.018	0.005	0.018	0.004
	L.S.D (0.05)	N.S.					
Linoleic (%)	D1	0.12	0.63	0.28	0.63	0.28	0.63
	D2	0.63	0.53	0.60	0.53	0.63	0.63
	D3	0.33	0.70	0.47	0.70	0.37	0.60
	L.S.D (0.05)	N.S.					
Palmitic (%)	D1	9.65	8.17	7.95	8.17	7.28	6.57
	D2	8.17	5.17	4.37	5.17	6.57	5.90
	D3	2.24	5.77	5.17	5.77	3.77	5.03
	L.S.D (0.05)	N.S.					
Stearic (%)	D1	3.78	9.48	4.88	9.48	4.42	6.80
	D2	9.48	5.27	3.90	5.27	6.80	6.33
	D3	1.73	5.43	5.70	5.43	4.17	4.37
	L.S.D (0.05)	N.S.					
Myristic (%)	D1	0.03	0.07	0.05	0.07	0.04	0.06
	D2	0.07	0.05	0.05	0.05	0.06	0.06
	D3	0.03	0.06	0.05	0.06	0.03	0.06
	L.S.D (0.05)	N.S.					

Table (6) Effect of Dates and Quantity of Potassium Application on Yield traits during Spring Season 2012

K treatments	No. seeds Seed.head ⁻¹	1000-seed weight (gm)	Plant yield (gm)	Total seed yield (ton.ha ⁻¹)	Oil content (%)	Oil yield (ton.ha ⁻¹)
K0	709	37.0	39.6	2.64	15.18	0.424
K1	860	31.8	46.0	3.07	14.52	0.446
K2	668	79.2	38.1	2.54	13.98	0.365
K3	821	84.2	55.9	3.72	12.84	0.483
K4	597	44.6	30.2	2.01	16.12	0.332
K5	716	31.4	26.1	1.74	14.50	0.253
L.S.D 0.05	121.6	24.34	15.42	1.028	N.S.	N.S.
Dates						
T1	721	60.5	41.8	2.78	13.19	0.362
T2	685	41.9	32.1	2.14	13.73	0.297
T3	779	51.7	44.1	2.94	16.65	0.492
L.S.D 0.05	N.S.	N.S.	N.S.	N.S.	1.772	0.1255

Table (7) Effect of interactions between Dates and Potassium Quantities on Yield traits during Spring Season.

Traits	Dates	Potassium Quantities					
		K0	K1	K2	K3	K4	K5
Number of Seeds	D1	636	870	586	900	513	823
	D2	690	817	589	783	553	678
	D3	801	894	830	778	727	647
	L.S.D (0.05)	N.S.					
1000 Seeds Weight (gm)	D1	54.8	29.5	95.0	97.2	46.3	40.0
	D2	30.2	30.4	68.1	62.0	33.7	27.0
	D3	25.9	35.5	74.4	93.3	53.9	27.3
	L.S.D (0.05)	N.S.					
Seed Yield of plant (gm)	D1	34.7	48.7	38.5	72.9	29.7	26.1
	D2	34.7	42.7	28.6	33.4	29.0	24.0
	D3	49.3	46.6	47.2	61.3	31.9	28.2
	L.S.D (0.05)	N.S.					
Total seed Yield (ton.ha ⁻¹)	D1	2.31	3.25	2.56	4.86	1.98	1.74
	D2	2.32	2.84	1.91	2.23	1.93	1.60
	D3	3.29	3.10	3.15	4.09	2.13	1.88
	L.S.D (0.05)	N.S.					
Oil percent %	D1	11.46	11.70	13.46	12.92	15.46	14.14
	D2	13.98	15.42	11.13	13.16	13.64	14.90
	D3	20.10	16.46	17.18	12.46	19.25	14.46
	L.S.D (0.05)	N.S.					
Oil yield	D1	0.262	0.383	0.342	0.628	0.309	0.246
	D2	0.332	0.443	0.214	0.292	0.263	0.240
	D3	0.676	0.446	0.540	0.528	0.425	0.272
	L.S.D (0.05)	N.S.					

Table (8) Effect of Dates and Quantity of Potassium Application on Fatty acids (%) during Spring Season 2012

K treatments	Fatty acids contents (%)				
	Oleic (%)	Linoleic (%)	Palmitic (%)	Stearic (%)	Myristic (%)
K0	0.025	0.36	6.68	5.00	0.04
K1	0.028	0.62	6.34	6.61	0.06
K2	0.050	0.36	6.46	4.78	0.04
K3	0.023	0.60	6.37	6.73	0.06
K4	0.026	0.36	6.68	5.00	0.04
K5	0.024	0.60	6.14	6.61	0.06
L.S.D 0.05	N.S.	0.175	N.S.	N.S.	N.S.
Dates					
T1	0.037	0.37	8.90	6.57	0.05
T2	0.034	0.56	6.50	7.21	0.06
T3	0.016	0.52	3.95	3.58	0.04
L.S.D 0.05	N.S.	0.123	1.161	2.45	N.S.

Table (9) Effect of interactions between Dates and Potassium Quantities on Fatty Acids (%) during Spring season 2012

Fatty acids	Dates	Potassium Quantities					
		K0	K1	K2	K3	K4	K5
Oleic (%)	D1	0.030	0.045	0.030	0.045	0.030	0.045
	D2	0.042	0.026	0.052	0.019	0.045	0.022
	D3	0.002	0.015	0.069	0.005	0.002	0.005
	L.S.D (0.05)	N.S.					
Linoleic (%)	D1	0.12	0.62	0.12	0.62	0.12	0.62
	D2	0.63	0.53	0.63	0.47	0.63	0.47
	D3	0.33	0.70	0.33	0.70	0.33	0.70
	L.S.D (0.05)	N.S.					

	L.S.D (0.05)	0.302					
Palmitic (%)	D1	9.65	8.10	9.65	8.17	9.65	8.17
	D2	8.17	5.17	7.83	5.17	8.17	4.50
	D3	2.24	5.77	1.90	5.77	2.24	5.77
	L.S.D (0.05)	2.845					
Stearic (%)	D1	3.78	9.48	3.78	9.48	3.78	9.14
	D2	9.48	4.93	8.81	5.27	9.48	5.27
	D3	1.73	5.43	1.73	5.43	1.73	5.43
	L.S.D (0.05)	N.S.					
Myristic (%)	D1	0.03	0.07	0.03	0.07	0.03	0.07
	D2	0.07	0.05	0.07	0.05	0.07	0.05
	D3	0.03	0.06	0.02	0.06	0.03	0.06
	L.S.D (0.05)	0.040					

IV. Conclusion

Results indicated that application of potassium at second date T_2 increased Seed number, 1000 seed weight, Plant yield and total seed yield, whereas Oil content and yield were increased with date of potassium application especially at third date T_3 . According to these results, it may recommended that fertilization of sunflower crop with K_3 treatment at third date T_3 for obtaining higher seed yield during both seasons. Also fertilization of potassium at third date when the aim is increasing content and yield of oil.

Acknowledgements

References

- [1] Murphy, W.M. 1978. Effects of planting date on seed oil and forage yield of irrigated sunflower.
- [2] Syed, A. S., M. Shahid, A. Jan and S. Noor-ud-din 2000. Effect of various Levels of Nitrogen, Phosphorus and Potassium (NPK) on growth, Yield and yield components of Sunflower. Pakistan Journal of Biological Sciences. 2000, Vol. 3, (2) pp:338-339.
- [3] Abdel-Motagally F. M. F. and E. A. Osman 2010. Effect of Nitrogen and Potassium Fertilization combination on productivity of two Sunflower cultivars under east of El-ewinatecondition. American - Eurasian Journal Agric. & Environ. Sci.8 (4). 2010, pp. 397-401
- [4] Chaudhry, A. U., and M. Mushtaq. 1999. Optimization of potassium in Sunflower. Pakistan Journal of Biological Sciences. 2 (3), pp. 887-888.
- [5] Alzobay, M.A. S. and J. H. Hamza. 2003. Effect levels of phosphorous and potash fertilizer on growth and yield sunflower. Morphological and Quality Traits. Iraqi journal of Agricultural Sciences. Pp 111-120, Volume 6 (34).
- [6] Abodhahy, Y.M. and M.A. Alyones. 1988. Plant Nutrition guide. Ministry of higher education and scientific research. Baghdad university.
- [7] Page, A. L. and et.al. 1982. Methods of soil analysis. Part (2) Agronomy No. 9. Madison. USA: s.n., 1982.
- [8] Alaloosy, Y.A.M. 2002. Effect of interaction between application of K, N and P fertilizer on growth and yield of sunflower. Iraqi journal of Agricultural Sciences, Vol. 3(33).
- [9] Lahmood, A.M. 1999. Effect of different dates of foliar fertilization on increasing percent of fertilization and yield seeds of two cultivars of sunflower. Technical college/Almusaib. Al-qadesyah journal. 2006, Vol. 11(1), Pp 1-13.
- [10] Salama, A. (1987). Yield and oil quality of sunflowers as affected by fertilizers and growth regulators. (36), pp. 191-202.
- [11] Mekki, B. (1999). Yield, oil and fatty acids content as affected by water deficit and potassium fertilization in two sunflower cultivars. Egypt. J. Agron. (21), pp. 67-85.
- [12] Ahmad, R., M. Saeed, E. Ullah and T. Mahmood 1999. Effect of Potassium on protein, Oil and Fatty acids contents in Two Autumn planted Sunflower Hybrids. International Journal of Agriculture & Biology. 1999, Vol. 1, 4, pp. 325-327.
- [13] Adnan, Y.Y. 2011. Effect of foliar fertilization ALGATON on growth traits, seed and oil yield of sunflower *Helianthus annuus* L. Iraq flower c.v. Tekreet journal of Agricultural Sciences, vol. 11(3), Pp 102-109.
- [14] Aldalayme, H.Y. and Y.A. Alaloosy. 2001. Effect of compound fertilizer (N, P, K) using foliar and soil application on yield and yield components of sunflower plant. Iraqi journal of Agricultural Sciences, Vol. 32(4), Pp 55-58.
- [15] Grove, J. H. and M. E. Sumner 1982. Yield and leaf composition of sunflower in relation to N, P, K, and lime treatments. Journal Nutrient Cycling in Agroecosystems. 1982, Vol. 3,(4), pp. 367-378.