

## Interdependency between Rainfall and Temperature using Correlation Analysis in the Barishal District of Bangladesh

Mst. Tania Islam<sup>1</sup>, Md. Zakaria<sup>2</sup>

<sup>1</sup>(Oceanographic Data Center, Bangladesh Oceanographic Research Institute, Cox's Bazar, Bangladesh)

<sup>2</sup>(Geological Oceanography Division, Bangladesh Oceanographic Research Institute, Cox's Bazar, Bangladesh)

**Abstract:** The paper tested the null hypothesis that there is no statistically significant correlation between rainfall and temperature for 68 years (1949 – 2017) of data. The analyses involved bivariate correlation and linear regression. Annual rainfall range was 1277 mm – 3390 mm, maximum temperature was 32.2 °C – 43.6 °C and minimum temperature was 27.8 °C -5 °C. The highest mean maximum temperature was 34.50 °C and lowest mean maximum temperature was 32.00 °C. The lowest mean minimum temperature occurred in 1973 (13 °C) and the highest was recorded in 1965 (27.50 °C) where highest mean rainfall was recorded in 1960 (273 mm) and lowest in 1963 (100 mm). The results indicate that for Barishal district, there are negative correlations between rainfall and Maximum temperature for the months from February to May and in August and December. There is no statistically significant correlation between rainfall and maximum temperature. Negative correlations found between rainfall and minimum temperature from September to December, February and June also. There is no statistically significant correlation between rainfall and minimum temperature. The correlation between rainfall and temperature revealed that rainfall is not dependent on the variability of the temperature in Barisal region of Bangladesh.

**Keywords:** Barisal District; Rainfall; Maximum Temperature; Minimum Temperature; Correlation Analysis

Date of Submission: 02-10-2019

Date of Acceptance: 18-10-2019

### I. Introduction

The climate of Bangladesh is changing and it is becoming more unpredictable every year [1]. Global warming induced changes in temperature and rainfall are already evident in many parts of the world, as well as in our country [2]. Hazards like floods, droughts, cyclones and others, which are aggravated by climate change and its variability being experienced more frequently in Bangladesh than ever before [3].

Therefore, Bangladesh has given special emphasis on research works on location specific climate change and its impact analyses, particularly at district level for crops, livestock, and forestry [4]. In Bangladesh the interest in climate-natural environment nexus is very much sustained. "Rainfall patterns show great fluctuations over the years and across vegetation zones. However there is gradual decrease in rainfall distribution in all parts of the country and this will affect agricultural production levels. Generally global temperatures are observed to be changing. The trend shows rising temperatures. The general outlook is that temperatures will continue to rise [5, 6]. According to Owusu-Sekyere [7] the onset of rainfall is presently starting late than previously and that annual rainfall is decreasing and temperature increasing. In fact, an accurate joint analysis of precipitation and temperature is more difficult to be carried out because of the possible interdependence between them [8]. Nevertheless, Rajeevan et al. [9] found that temperature and rainfall in India were positively correlated during January and May, but negatively correlated during July. Huang et al. [10] showed a negative correlation between rainfall and temperature in the Yellow River basin of Chin. In addition, utilization of water by crops (e.g. maize) has been consistent over the years; various crops show yield decline as a result of the decreasing total annual rainfall or the increasing mean temperatures or both. Actually, Bangladesh crop farming is mainly rain-fed, crop productivity is highly dependent on rainfall which varies greatly over short distances, hence, it is important to use rainfall data from the site of interest [11]. In this regard, the present paper examines local rainfall and temperature data in Barisal region for a period of 68 years. This analyses that follow show the correlation and regression between rainfall and temperature.

### II. Materials And Methods

To take the interdependence between rainfall and temperature into account, multivariate probability simulation is needed. Traditionally multivariate probability density functions, however, are generally limited to the multivariate normal distribution or mixtures of it [12]. This approach also allows for the comparisons of data series of different time periods and lengths [13].

The correlation analysis measures the degree of interdependence between two or more variables. It cannot prove a causal relationship, a relationship of cause and effect between variables. Interdependence can however be functional. By functional relationship we understand the relationship that can be expressed by a formula or by what mathematicians call a mathematics law, such as linear relationship formula. The fact that the two variables tend to be related, meaning that one of them increased levels tend to be accompanied by an increase of the second and vice versa, it doesn't turn out that the first has a direct influence on the second or vice versa.

Determination of relationship between two variables raises the question: how close, how intense are these relationships and, consequently, how much can vary estimates or predictions made on the basis of regression analysis. As the average cannot be properly interpreted without a measure of the dispersion or variability of the data from which it resulted (and the most common measure is the average square deviation or standard deviation), so estimates or predictions resulting from the regression analysis require finding a measure of their variability. We will consider as measures of estimations variability based on.

**Study Area:** The study area is situated in the most southern part of Bangladesh where Ganges-Brahmaputra-Meghna River system meets the Bay of Bengal. The study is only based on meteorological data (Rainfall, Maximum Temperature and Minimum Temperature) for the Barishal District of Bangladesh.



**Figure 1:** Study area and location of the station in Barisal District of Bangladesh.

**Data Collection and Preliminary Analysis:** Monthly maximum temperature, minimum temperature and rainfall data of Barishal district from 1949 to 2017 was obtained from the Bangladesh Meteorological Department (BMD). Data has been analyzed with Statistical Package for Social Scientists (SPSS 25 version) specifically using bivariate correlation and linear regression. Graphs were produced with Microsoft Excel.

### III. Results and Discussion

The statistical approach here used to explore the relationships between climatic data series which are not perfectly similar, such as monthly rainfall and temperature, is the correlative analysis applied to the standardized anomalies [14]. As first step, variability of average maximum temperature, minimum temperature and rainfall is listed in table. Then correlation between maximum temperature and rainfall, minimum temperature and rainfall has been delineated using SPSS.

**Maximum Temperature:** From the Table no 1, we see that the max. Temperature usually reaches its peak in April and its bottom in January; the average max. temperature is also peak in April and its bottom in January. The variability of average temperature in February and April is though relatively large. Some descriptive temperature statistics are listed in Table 1.

**Table no 1:** Descriptive statistics for monthly average maximum temperature from 1949 to 2017 (°C).

Max. Temp.	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sep.	Oct.	Nov.	Dec.
Maximum	31.7	35.6	38.4	43.6	38.6	39.6	35.5	36.7	37.7	35.6	34	31.7
Minimum	26.7	29.2	33.4	32.5	33.5	32.2	31.9	31.4	32.2	32.3	29.4	26.6
Mean	28.98	32.14	35.69	36.36	35.95	35.23	33.75	34.15	34.42	34.06	32.09	29.19
Standard deviation	1.065	1.321	1.075	1.642	0.991	1.368	0.890	1.082	0.929	0.769	1.124	1.049
Coefficient of variation	3.676	4.111	3.012	4.516	2.758	3.884	2.638	3.168	2.698	2.259	3.501	3.595

**Minimum Temperature:** From the Table No 2, we see that the min. temperature usually reaches its peak in August and its bottom in January; the average min. temperature is also peak in August and its bottom in January. The variability of average min. temperature in December and January is though relatively large.

**Table no 2:** Descriptive statistics for monthly average minimum temperature from 1949 to 2017 (°C).

Min. Temp.	Jan.	Feb.	March	April	May	June	July	Aug.	Sep.	Oct.	Nov.	Dec.
Maximum	12.6	14.4	20	23.5	26	26.2	25.4	25.7	25.6	23.5	18.7	15.6
Minimum	6.4	7.2	12	15.8	17	19.2	17.4	21.6	21.5	12.7	11.9	7.5
Mean	9.388	11.081	15.558	19.734	20.931	23.022	24.036	24.18	23.760	20.620	15.427	11.052
Standard deviation	1.606	1.685	1.889	1.445	1.343	1.138	1.227	0.888	0.929	1.661	1.520	1.587
Coefficient of Variation	17.117	15.203	12.142	7.322	6.418	4.942	5.104	3.672	3.911	8.056	9.854	14.359

**Rainfall:** From the Table No 3 we can say that June to September, the average monthly total rainfall is relatively high.

**Table no 3:** Descriptive statistics for monthly average Rainfall from 1949 to 2017 (mm).

Rainfall	Jan.	Feb.	March	April	May	June	July	Aug.	Sep.	Oct.	Nov.	Dec.
Maximum	148	98	273	412	701	1025	785	823	1067	497	274	197
Minimum	0	0	0	0	22	139	185	138	90	6	0	0
Mean	8.896	19.657	46.015	96.939	208.179	408.642	424.761	356.612	301.500	189.149	42.836	8.776
Standard deviation	23.122	24.265	62.419	85.603	123.278	175.814	141.945	142.239	160.344	120.749	63.421	26.754
Coefficient of Variation	259.932	123.445	135.647	88.306	59.217	43.024	33.418	39.886	53.182	63.838	148.055	304.851

**The Relationship between Rainfall and Maximum Temperature:** The physical rationale behind the relationship between rainfall and Maximum Temperature is that rainfall may affect soil moisture which may in turn affect surface temperature by controlling the partitioning between the sensible and latent heat fluxes. Because the sample data is non-Gaussian distributed and skewed, the Kendall correlation coefficient is employed to calculate the correlation between monthly rainfall and maximum temperature. It is found that there are negative correlations between rainfall and maximum temperature from February to May and in August and December (at the 10% confidence level) (Table 4). In the Table No 4, we see that p value is greater than level of significance. So we can not reject the null hypothesis. Moreover, we can say that there is no statistically significant correlation between rainfall and maximum temperature.


**Table No 4:** Correlation analysis for monthly max temperature and rainfall from 1949 to 2017.

	Jan.	Feb.	March	April	May	June	July	Aug.	Sep.	Oct.	Nov.	Dec.
Kendall correlation coefficients	.180	-.106	-.051	-.083	-.018	.036	.088	-.147	.034	.012	.025	-.018
P value	.059	.234	.570	.352	.839	.680	.315	.090	.704	.896	.783	.854

**The Relationship between Rainfall and Minimum Temperature:** The physical rationale behind the relationship between rainfall and Minimum Temperature is that rainfall may affect soil moisture which may in turn affect surface temperature by controlling the partitioning between the sensible and latent heat fluxes. Because the sample data is non-Gaussian distributed and skewed, the Kendall correlation coefficient is employed to calculate the correlation between monthly rainfall and minimum temperature. It is found that there are negative correlations between rainfall and minimum temperature from September to December and in February and June also (at the 10% confidence level). In the Table 5, we see that p value is greater than level of significance. So we can not reject the null hypothesis. Moreover, we can say that there is no statistically significant correlation between rainfall and .minimum temperature.


**Table No 5:** Correlation analysis for monthly min temperature and rainfall from 1949 to 2017

	Jan.	Feb.	March	April	May	June	July	Aug.	Sep.	Oct.	Nov.	Dec.
Kendall correlation coefficients	.056	-.051	.020	.022	.003	-.026	.137	.034	-.081	-.014	-.095	-.129
P value	.558	.566	.827	.802	.977	.771	.115	.701	.360	.873	.287	.188




**Figure 2:** Correlation plot between Rainfall and Max temperature

From SPSS output shown in figure 2, we see that correlation value of maximum temperature and rainfall is  $-.095$ . It can be said that, there is no relationship between maximum temperature and rainfall. No correlation means that the two sets of data are not related at all. In other words, it means one set of data does not increase or decrease with the other.



**Figure 3:** Correlation plot between Rainfall and Min temperature

From SPSS output shown in figure 3, we see that correlation value of minimum temperature and rainfall is  $-0.098$ . It also can be said that, there is no relationship between minimum temperature and rainfall. No correlation means that the two sets of data are not related at all. In other words, it means one set of data does not increase or decrease with the other.



**Figure 4:** Maximum temperature graph for 1949-2017

Figure 4 shows the maximum temperature for 68 years (1949-2017). The trend line indicates slightly increasing temperature. The highest mean maximum temperature was  $34.50$  ( $^{\circ}\text{C}$ ) and lowest mean maximum temperature was  $32.00$  (deg.cel).The mean maximum temperature is generally favorable for the growing of any of the major crops.




Figure 5: Minimum temperature graph for 1949-2017

Figure 5 reveals an ascending trend line for minimum temperature for 1949-2017. From the 68 years of data, the lowest mean minimum temperature occurred in 1973 (13°C) and the highest was recorded in 1965 (27.50°C).




Figure 6: Rainfall graph for 1949-2017

Figure 6 portrays trend of rainfall from 1949 to 2017. From the graph we can say that, highest mean rainfall was recorded in 1960 (273mm) and lowest in 1963 (100mm).

#### **IV. Conclusions**

This paper presents a correlation analysis of temperature and rainfall, which are of utmost importance for agricultural production especially in the context of climate change. The main purpose of this paper is to present interdependence of rainfall and temperature in Barishal region, Bangladesh. The study is only based on meteorological data for a single region. A potentially valuable extension of this research is to connect the analysis with crop production planning and agricultural economics. If the relationship among temperature, rainfall, and crop yield can be determined, then it could be used in developing risk reducing strategies for farmers, something which will become increasingly important in the face of climate change. This is the focus of our ongoing research. From the analysis of 1949-2017 years of data, it has been found that annual rainfall range was 1277 mm – 3390 mm, maximum temperature was 32.2 °C – 43.6 °C and minimum temperature was 27.8 °C -5 °C. The highest mean maximum temperature was 34.50 °C and lowest mean maximum temperature was 32.00 °C. The lowest mean minimum temperature occurred in 1973 (13 °C) and the highest was recorded in 1965 (27.50 °C) where highest mean rainfall was recorded in 1960 (273 mm) and lowest in 1963 (100 mm). Finally, The correlation between rainfall and temperature revealed that rainfall is not dependent on the variability of the temperature in Barishal region of Bangladesh.

#### **References**

- [1]. World Bank. Bangladesh: Climate Change and Sustainable Development Report No. 21104 RD. World Bank Office, Dhaka. South Asia Rural Development Team. 2000
- [2]. IPCC, "Summary for Policymakers: A Report of Working Group I of the Intergovernmental Panel on Climate Change." 2001. URL: <http://www.ipcc.ch/pub/spm22-01.pdf>.
- [3]. IPCC, Climate Change 2007: Impacts, Adaptations, and Vulnerability, 2007; Contribution of Working
- [4]. Ahmed, A.U. and M.A. "Development of Climate Change Scenarios with General Circulation. 1999.
- [5]. IPCC, Climate Change 2007: Impacts, Adaptation and Vulnerability, WMO, Geneva, 2007; pp. 1-23.
- [6]. Environmental Protection Agency, Ghana State of the Environment Report 2004, Environmental Protection Agency, Accra, 2005; pp. 1-152.
- [7]. J. D. Owusu-Sekyere, J. Andoh, and K. Nyarko, Climate Change and Crop Production in the Mfantseman Area of Ghana. J. Appl. Environ. Biol. Sci. 2011; 1, 134-141.
- [8]. Tanarhte, M.; Hadjinicolaou, P.; Lelieveld, J. Intercomparison of temperature and precipitation data sets based on observations in the Mediterranean and the Middle East. J. Geophys. Res. 2012; 117.
- [9]. Rajeevan, M.; Pai, D.S.; Thapliyal, V. Spatial and temporal relationships between global land surface air temperature anomalies and Indian summer monsoon rainfall. Meteorol. Atmos. Phys. 1998; 66, 157–171
- [10]. Huang, Y.; Cai, J.; Yin, H.; Cai, M. Correlation of precipitation to temperature variation in the Huanghe River (Yellow River) basin during 1957–2006. J. Hydrol. 2009, 372, 1–8.
- [11]. D.K. Asare, and H.M. Amoatey, Potential of Crop Models for Improving and Sustaining Crop Production in Ghana. Ghana Jnl Agric. Sci. 2001; 34, 127-133.
- [12]. D. S. Wilks, Statistical Methods in the Atmospheric Sciences, Academic Press, 2nd edition, 2005.
- [13]. Rodrigo, F.S. On the covariability of seasonal temperature and precipitation in Spain, 1956–2005. Int. J. Climatol. 2014; 35, 3362–3370
- [14]. Wilks, D.S. Statistical Methods in the Atmospheric Sciences, 2nd ed.; International Geophysics Series; Academic Press: Cambridge, MA, USA, 1995.

Mst. Tania Islam. "Interdependency between Rainfall and Temperature using Correlation Analysis in the Barisal District of Bangladesh." IOSR Journal of Mathematics (IOSR-JM) 15.5 (2019): 49-55.