

A Review on Physiological, Biochemical and Biotechnological Applications of Mushroom

Sanjay Mishra^{1,*} and R.B. Singh²

¹School of Biotechnology, IFTM University, Lodhipur Rajput, Delhi Road (NH-24), Moradabad 244 102, U.P., India, ²Halberg Hospital and Research Institute, Civil Lines, Moradabad 244 001, U.P., India

*Corresponding Author: Email: sanjaymishra@iftmuniversity.ac.in

Abstract: Mushrooms are a manifestation of a common saying, 'Medicines and foods have a common origin', in constituting both a nutritionally functional food and a source of physiologically beneficial medicine. The review article covers the data obtained from the broad-range studies focused on physiological and biochemical worth of varieties as well as mode of nutrition of mushroom and may provide an overview for physiologists, biochemists and biotechnologists to propagate their research on still obscure nutraceutical significance of edible mushrooms.

Keywords: L-carnitine, Lipid peroxidation, Lipid profile, Liver function, Mushroom, Mushroom extract, Swiss Albino rat,

Introduction

In fact, aging is associated with biochemical and structural alterations which are thought to result in motor and cognitive impairments and in increased susceptibility to neurodegenerative diseases¹⁻⁴. The free radical theory of aging proposed that aging is due to the accumulation of unrepaired damage from free radical attack on cellular components. Modern approaches propose that aging is caused by a shift in the balance between the pro-oxidative and anti-oxidative processes in the direction of the pro-oxidative state^{1,2,5-7}. L-carnitine, a nutrient normally synthesized from methionine and lysine in the liver and kidney. L-carnitine transports long-chain fatty acids (LCFA) across the mitochondrial membrane where they undergo beta-oxidation to produce energy. Carnitine deficiency decreases LCFA availability for oxidation, thereby resulting in LCFA accumulation in the cytosol, and decreased ketone and energy production. Other L-carnitine functions include the maintenance of adequate free coenzyme-A required for various metabolic pathways, the protection of cells against toxic accumulation of acyl-coenzyme-A compounds by shuttling acyl groups out of the mitochondria, and the storage and transport of energy⁸. Also, L-carnitine supports the immune system and enhances the antioxidant system⁹.

Mushrooms are a manifestation of a common saying, 'Medicines and foods have a common origin', in constituting both a nutritionally functional food and a source of physiologically beneficial medicine. Many centuries ago, medicinal properties of mushrooms have been recognized in China, Korea and Japan. Although from ancient times, mushrooms have been treated as a special kind of nutraceutical, they have received a remarkable interest in recent decades. Major medicinal properties attributed to mushrooms include anticancer activity, antibiotic activity, antiviral activity, immune response-stimulating effects, anti-hypersensitive and blood lipid lowering effects¹⁰⁻¹². Mushroom is known to have high amounts of proteins, carbohydrates and fibers and low fat contents¹³. Furthermore, mushroom had significant levels of vitamins, namely thiamine, riboflavin, ascorbic acid and vitamin D₂, as well as minerals¹⁴. Mushroom species had been shown to possess antioxidant capacity in *in-vitro* systems¹⁵⁻¹⁸. The mushroom *Pleurotus species* (*P. ostreatus*, *P. sajor-caju*, *P. florida*) were reported to have hypocholesterolemic activity in experimental rats¹⁸⁻²⁰. It has been reported that the L-carnitine concentration in mushroom ranged from 130 to 533 mg/kg dried mushroom²¹. The free L-carnitine concentration in mushroom ranged from 75 to 385 mg/kg dried mushroom, which represented 70 ± 10% of total carnitine content. The present report is the compilation of overall studies focusing at physiological as well as biochemical significance of mushroom projecting the nutraceutical visage for researchers to propagate the research in relevant thrust area.

Alteration in lipid physiology and biochemistry of mammalian system upon mushroom feeding

The effect of 15% dried mushroom, 450 mg mushroom extract and L-carnitine on total lipid, triglyceride and total cholesterol has been well described in studies accomplished previously^{21,22}. Total lipid content significantly ($P \leq 0.05$) reduced in albino rats supplemented with mushroom and L-carnitine. The reduction in the total lipids ranged from 7.06 to 14.39%. There was no significant ($P > 0.05$) difference in total lipid between rats supplemented with 400 mg L-carnitine and those supplemented with 15% dried mushroom. Albino rats supplemented with 800 mg L-carnitine had a higher ($P \leq 0.05$) total lipid content compared to those

supplemented with 450 mg mushroom extract. Diet supplemented with mushroom and L-carnitine resulted in a significant ($P \leq 0.05$) decrease in triglyceride and total cholesterol level. Triglyceride was observed to reduce by 31.28–43.72%. However, total cholesterol reduced by 15.92–28.45%. Supplementation with 450 mg mushroom extract and 800 mg L-carnitine were more ($P \leq 0.05$) effective in reducing triglyceride and total cholesterol than those supplemented with 15% dried mushroom and 400 mg L-carnitine. On the other hand, supplementation with 450 mg mushroom extract and 800 mg L-carnitine were similar ($P > 0.05$) in reducing triglyceride and total cholesterol levels. Supplementation with 15% dried mushroom and 400 mg L-carnitine were also similar ($P > 0.05$) in reducing triglyceride and total cholesterol. It has been observed that rats fed a semisynthetic diet containing 0.3% cholesterol and supplemented with 5% dried whole oyster mushroom had reduced serum and liver cholesterol levels by 34 and 58%, respectively²¹. Panchamoorthy and Carani²² reported that treated rats with L-carnitine caused a significant reduced in TG as compared to untreated rats. L-carnitine is known to promote the transport of cytosolic long-chain fatty acids into the mitochondrial matrix for β -oxidation, thereby providing mitochondrial energy^{23, 24}. L-carnitine may lower plasma TG by increasing the utilization and/or oxidation of fatty acids for energy or possibly by altering very low-density lipoprotein synthesis²⁵.

The data gathered so far²²⁻²⁵ reflect that the high density lipoprotein in rats was not affected ($P > 0.05$) as a consequence of the supplementation with 15% dried mushroom and 400 mg L-carnitine. However, rats supplemented with 450 mg mushroom extract and 800 mg L-carnitine had a higher ($P \leq 0.05$) high density lipoprotein compared to those of the control sets. High density lipoprotein was monitored to enhance in these albino rats by 24.11–30.44%. Low density lipoprotein ($P \leq 0.05$) reduced in albino rats supplemented with mushroom and L-carnitine by 30.36–55.76%. Supplementation of rats with 450 mg mushroom extract and 800 mg L-carnitine were more ($P \leq 0.05$) effective in lowering low density lipoprotein than those supplemented with 15% dried mushroom and 400 mg L-carnitine. On the other hand, supplementation of rats with 450 mg mushroom extract and 800 mg L-carnitine were similar ($P > 0.05$) in decreasing low density lipoprotein. Supplementation of rats with 15% dried mushroom and 300 mg L-carnitine were also similar ($P > 0.05$) in lowering low density lipoprotein. Very low density lipoprotein in rats was ($P \leq 0.05$) reduced by the supplementation with mushroom and L-carnitine. Very low density lipoprotein was reduced in these rats by 32.33–42.21%. Supplementation of rats with 450 mg mushroom extract and 800 mg L-carnitine were more ($P \leq 0.05$) effective in reducing very low density lipoprotein than those supplemented with 15% dried mushroom and 400 mg L-carnitine. Diet supplemented with 450 mg mushroom extract and 800 mg L-carnitine did not significantly ($P > 0.05$) differ in their effect on very low density lipoprotein. Besides, no significant ($P > 0.05$) difference was observed in very low density lipoprotein between albino rats supplemented with 15% dried mushroom and those supplemented with 400 mg L-carnitine. These results are in agreement with those reported earlier^{26, 27} highlighting that L-carnitine well stabilizes the level of lipids peroxidation, decreases concentration of total lipids, triglycerides, total cholesterol, phospholipids, and lipoproteins of low and very low density, in the Swiss albino rats' blood sera.

Alteration in the physiological and biochemical level of major enzymes concerning with liver function of mammalian system upon feeding mushroom supplemented with L-carnitine

The aspartate amino transferase (AST) enzyme in the mammalian system was observed to considerably reduce as a consequence of the supplementation of diet with mushroom and L-carnitine. Mushroom reduced AST enzyme by 38.64–41.46%. However, L-carnitine reduced it by 24.58–42.80%. Swiss albino rats supplemented with 400 mg L-carnitine showed a higher ($P \leq 0.05$) AST enzyme compared to those supplemented with mushroom and 800 mg L-carnitine. Diet supplemented with 450 mg mushroom extract and 800 mg L-carnitine were not significantly ($P > 0.05$) differed in their impact on AST enzyme. Further, diet supplemented with mushroom and L-carnitine had a lower ($P \leq 0.05$) alanine amino transferase (ALT) enzyme compared to that of the control sets. Mushroom and L-carnitine reduced ALT enzyme by 36.59–45.61% and 22.40–36.99%, respectively. Diet supplemented with 15% dried mushroom, 450 mg mushroom extract and 800 mg L-carnitine appeared to be more effective ($P > 0.05$) in decreasing ALT enzyme compared to those supplemented with 400 mg L-carnitine. No significant ($P > 0.05$) difference was found in ALT enzyme among rats supplemented with 15% dried mushroom, 450 mg mushroom extract and those supplemented with 800 mg L-carnitine. The alkaline phosphatase (ALP) enzyme in rats was observed to significantly ($P \leq 0.05$) reduce by the supplementation with mushroom and L-carnitine. Mushroom reduced ALP enzyme by 22.19–32.71%. However, L-carnitine reduced it by 22.14–49.26%. The Diet supplemented with 400 mg L-carnitine had a higher ($P \leq 0.05$) ALP enzyme compared to those supplemented with 800 mg L-carnitine. Diet supplemented with 15% dried mushroom had a higher ($P \leq 0.05$) ALP enzyme compared to those supplemented with 450 mg mushroom extract. The diet supplemented with 15% dried mushroom and 400 mg L-carnitine were not significantly ($P > 0.05$) differed in their impact on ALP enzyme. L-carnitine and mushroom restores the changes of ALT, AST and ALP activities due to their antioxidant effects and their ability to act as a radical scavenger, thereby protecting membrane permeability. It has been observed found that ALT and AST after ethanol intoxication

their activity increased by about 80%. L-carnitine partly prevented these changes. It was manifested by a statistically significant decrease in the activity of ALT and AST, by about 20% in comparison with the ethanol group^{27,28}.

The Data obtained from the studies^{27,28} reflect that the MDA ($P \leq 0.05$) got reduced by 11.92-33.79% in albino rats supplemented with diet containing mushroom and L-carnitine. Supplementation with 450 mg mushroom extract and 800 mg L-carnitine were more ($P \leq 0.05$) effective in decreasing MDA compared to those supplemented with 15% dried mushroom and 400 mg L-carnitine. On the other hand, supplementation with 450 mg mushroom extract and 800 mg L-carnitine were similar ($P > 0.05$) in reducing MDA. Supplementation of diets with 450 mg mushroom extract and 400 mg L-carnitine were also similar ($P > 0.05$) in reducing MDA. Rats supplemented with the diet containing 15% dried mushroom had higher ($P \leq 0.05$) MDA compared to those supplemented with 400 mg L-carnitine. It has earlier been reported that administration of L-carnitine to rats intoxicated with ethanol significantly protects lipids and proteins against oxidative modifications in the serum and liver. The level of MDA was decreased by about 30%, in the blood serum in comparison to the ethanol group²⁸.

Glutathione peroxidase (GSHPx) is known to perform a key role in co-coordinating the innate antioxidant defense mechanisms. It is involved in the maintenance of the normal structure and function of cells, probably by its redox and detoxification reactions^{1, 2, 29-31}. The GSHPx in rats was monitored to be significantly ($P \leq 0.05$) enhanced by the supplementation with mushroom and L-carnitine. Mushroom increased GSHPx by 58.43-85.50%. However, L-carnitine increased it by 60.15-129.69%. Rats supplemented with 450 mg L-carnitine and 15% dried mushroom had a lower ($P \leq 0.05$) GSHPx compared to those supplemented with 800 mg L-carnitine and 450 mg mushroom extract. Supplementation with 15% dried mushroom and 400 mg L-carnitine were not significantly ($P > 0.05$) differed in their effect on GSHPx. Supplementation of rats with 800 mg L-carnitine was more ($P \leq 0.05$) effective in increasing GSHPx compared to those supplemented with 400 mg L-carnitine, 15% dried mushroom and 450 mg mushroom extract. According to Augustyniak and Skrzydlewska²⁸ L-carnitine has been reported to cause a significant increase in the liver and blood serum GSH level by more than 20%. An increase in the levels of GSHPx in aged rats treated with mushroom extract as a source of antioxidant has also been recently reported.

The overall results outcome from the research studies³²⁻³⁹ highlight the effect of dried mushroom, mushroom extract and L-carnitine on food intake and body weight of Swiss albino rats. Either L-carnitine or mushroom significantly ($P \leq 0.05$) increased food intake and reduced body weight in rats. There was no significant ($P > 0.05$) variation in food intake between rats supplemented with L-carnitine and mushroom. Supplementation of rats with L-carnitine was more ($P \leq 0.05$) effective in reducing body weight than those supplemented with mushroom. Supplemented rats with 400 mg L-carnitine and 800 mg L-carnitine were not significantly ($P > 0.05$) distinct in their effect on body weight. Similar effect was monitored in rats supplemented with 15% dried mushroom and 450 mg mushroom extract. The rationale for L-carnitine supplementation as a weight-loss agent is based on the assumption that regular oral ingestion of the substance increases its intracellular concentration. This would trigger increased fat oxidation and gradual reduction of the body's fat reserves³²⁻³⁹.

Acknowledgements

The present piece of work was supported by a joint venture of Halberg Hospital and Research Institute, Civil Lines, Moradabad 244 001, U.P., India and School of Biotechnology, IFTM University, Lodhipur Rajput, Delhi Road (NH-24), Moradabad 244 001, U.P., India.

References

- [1] Sharma V, Kalim S, Srivastava MK, Nanda S, Mishra S. Oxidative stress and coxsakievirus infections act as mediators of beta cell damage: a review. *Sci Res Essay* 2009; 4 (2): 42-58
- [2] Nanda S, Mishra S, Varshney VP, Singh RB. A biotechnological approach to apoptosis of somatic and germ cells in living organisms. *The Open Nutraceuticals J* 2010; 3: 81-93.
- [3] Freo U, Pizzolato G, Dam M, Ori, C, Battistin L. A short review of cognitive and functional neuroimaging studies of cholinergic drugs: implications for therapeutic potentials. *J Neural Transm* 2002; 109: 857-70.
- [4] Terry J, Buccafusco J J. The cholinergic hypothesis of age and Alzheimer's disease-related cognitive deficits: recent challenges and their implications for novel drug development. *J Pharmacol Exp Ther* 2003; 306: 821-27.
- [5] Harman D. Free radical theory of aging. *Mutation Res* 1992; 275: 257-66.
- [6] Beckman B, Ames B.N. The free radical theory of aging matures. *Physiol Rev* 1998; 78: 547-81.
- [7] Cadenas E, Davies KJ. Mitochondrial free radical generation, oxidative stress and aging. *Free Radical Biol Med* 2000; 29: 222-30.
- [8] Catherine MC, Michael CS, Michael LC, *et al*. Carnitine supplementation in premature neonates: Effect on plasma and red blood cell total carnitine concentrations, nutrition parameters and morbidity. *Clinical Nutrition* 2006; 25, 886-96
- [9] Bremer J. The role of carnitine in cell metabolism. In: De Simone, C., Famularo, G. (Eds.), *Molecular Biology Intelligence Unit Carnitine Today* (Landes Bioscience Austin, TX, USA, International Copyright). Springer Verlag, Heidelberg, Germany; 1997; pp. 4-37.
- [10] Wasser SP, Weis AL. Medicinal properties of substances occurring in higher Basidiomycetes mushrooms: Current perspectives. *Int J Med Mushr* 1999; 1(1): 31-62.

- [11] Kaul TN. Biology and Conservation of mushrooms. New Delhi: Oxford & IBH Publishing Co. Pvt. Ltd 2001: pp. 117-45.
- [12] Yang JH, Lin HC, Mau JL. Antioxidant properties of several commercial mushrooms. *Food Chem* 2002; 77: 229-35.
- [13] Bárbara R, Rosário L, Paula B, *et al.* Comparative study of phytochemicals and antioxidant potential of wild edible mushroom caps and stipes. *Food Chem* 2008; 110: 47-56.
- [14] Mattila P, Suonpaa K, Piironen V. Functional properties of edible mushrooms. *Nutrition* 2000; 16: 694-96.
- [15] Ribeiro B, Rangel J, Valentão P, *et al.* Contents of carboxylic acids and two phenolics and antioxidant activity of dried portuguese wild edible mushrooms. *J Agric Food Chem* 2006; 54: 8530-37.
- [16] Diaz M, Lopez F, Hernandez F, Urbina JA. L-carnitine effects on chemical composition of plasma lipoproteins of rabbits fed with normal and high cholesterol diets, *Lipids* 2000; 35: 627-32.
- [17] Ribeiro B, Valentão P, Baptista P, *et al.* Phenolic compounds, organic acids profiles and antioxidative properties of beefsteak fungus (*Fistulina hepatica*). *Food Chem Toxicol*. doi:10.1016/j.fct.2007.03.015.
- [18] Khan MA, Tania M, Zhang D-Z, Chen H-C. Cordyceps mushroom: a potent anticancer nutraceutical. *The Open Nutraceuticals J* 2010; 3: 179-83.
- [19] Hossain S, Hashimoto M, Choudhury EK, *et al.* Dietary mushroom (*Pleurotus ostreatus*) ameliorates atherogenic lipid in hypercholesterolaemic rats. *Clin Exp Pharmacol Physiol* 2003; 30(7): 470-5.
- [20] Alam N, Amin R, Khan A, *et al.* Comparative effects of oyster mushrooms on lipid profile, liver and kidney function in hypercholesterolemic rats. *Mycobiology* 2009; 37(1): 37-42.
- [21] Bobek P, Ozdín L, Kuniak L. Mechanism of hypocholesterolemic effect of oyster mushroom (*Pleurotus ostreatus*) in rats: reduction of cholesterol absorption and increase of plasma cholesterol removal. *Z Ernährungswiss.* 1994; 33: 44-50
- [22] Panchamoorthy R, Carani, V. (2007). Fructose-induced hepatic gluconeogenesis: Effect of L-carnitine. *Life Sciences* 80 : 1176-83.
- [23] Eskandari HG, Burak Cimen MY, Tamer L, *et al.* Short term effects of l-carnitine on serum lipids in STZ-induced diabetic rats *Diabetes Research and Clinical Practice* 2004; 66: 129-32.
- [24] Tanaka Y, Sasaki R, Fukui F, Waki H, Kawabata T, Okazaki M, (2004). Acetyl-l-carnitine supplementation restores decreased tissue carnitine levels and impaired lipid metabolism in aged rats. *J Lipid Res.* 45:729-35.
- [25] Sidoriak NG, Volgin DV. Effect of L-carnitine on lipid peroxidation and lipid composition in blood serum in hemic hypoxia. *Ukr Biokhim Zh* 1996; 68(5): 54-8.
- [26] Lofgren I, Zern T, Herron K, *et al.* Weight loss associated with reduced intake of carbohydrate reduces the atherogenicity of LDL in premenopausal women. *Metabolism* 2005; 54(9):1133-41.
- [27] Seline KG, Johein H. The determination of L-carnitine in several food samples. *Food Chem* 2007; 105: 793 - 804.
- [28] Augustyniak A, Skrzydlewska E. L-Carnitine in the lipid and protein protection against ethanol-induced oxidative stress. *Alcohol* 2009; 43, 217-23.
- [29] American Institute of Nutrition (AIN), Purified diet for laboratory rodent. Final Report. ADHOC Writing Diet. *J. Nutr.* 1993; 123: 1939 - 51.
- [30] Black K. In: *Business Statistics for Contemporary Decision Making* (Fourth (Wiley Student Edition for India) ed.). Wiley-India. 2004: ISBN 9788126508099.
- [31] Gueeri H. Influence on prolonged ethanol intake on the level and turnover of alcohol and aldehyde dehydrogenase and glutathione. *Adv Exp Med Biol* 1995; 23: 133-34.
- [32] Bergmeyer HU, Harder M. A colorimetric method of the determination of serum glutamic oxaloacetic and glutamic pyruvic transaminase *Clin Biochem* 1986; 24: 481.
- [33] Kachmar JF, Moss DW. Enzymes, In: *Fundamentals of Clinical Chemistry* (edited by Tiez N). 1976; pp. 666-672, Philadelphia PA. W.B. Saunders Co.
- [34] Varley H, Gewenlock A, Bell M. *Practical clinical biochemistry*, Vol. 1, 5th ed. 1980; Pp.741:897. London; Williams Heinemen Medical books, Ltd
- [35] Hu ML. Measurement of protein thiol groups and glutathione in plasma. *Methods Enzymol* 1994; 233: 380-85.
- [36] Jentsch AM, Bachmann H, Furst P, Biesalski HK. Improved analysis of malonaldehyde in human body fluids. *Free Radic Biol Med* 1996; 20: 251-60.
- [37] Jayakumar T, Ramesh E, Geraldine P. Antioxidant activity of the oyster mushroom, *Pleurotus ostreatus*, on CCl₄-induced liver injury in rats. *Food Chem Toxicol* 2006; 44: 1989-96.
- [38] Akoma O, Onuoha SA, Ajiboye MO, Akoma AO, Alawoki AM. The nutritional and sensory quality characteristics of kunun-zaki produced with the addition of hydrolytic enzymes from malted rice (*Oryza sativa*). *J Food Technol Afr* 2002; 7: 24-6.
- [39] Akoma O, Jiya EA, Akumka DD, Mshelia E. Influence of malting on the nutritional characteristics of kanun-zaki. *AfrJ Biotechnol* 2006; 5: 996-1000.