

Antibiotic Sensitivity Pattern of Typhoid Fever- Experience from a Tertiary Care Hospital in Bangladesh

Ishrat Jahan¹, Jesmin Akter Mitu², Mehdi Pervez³, Sanjib Kundu⁴, Maruf Hossain⁵, Sheikh Anisul Haque⁶

¹Professor, Department of Pediatrics, Bangladesh Shishu Hospital and Institute, Dhaka, Bangladesh, Orcid Id: 0009-0005-7958-820X

²Junior Consultant, Department of Pediatrics and Neonatology, Bangladesh Specialized Hospital, Dhaka, Bangladesh, Orcid Id: 0000-0001-8155-3026

³Assistant Professor, Pediatrics (in situ), 250 Bed General Hospital, Barguna, Bangladesh, Orcid Id: 0000-0001-9863-0550

⁴Senior Medical Officer, Neonatal Intensive Care Unit, Bangladesh Specialized Hospital, Dhaka, Bangladesh,

⁵Register, Department of Pediatrics and Neonatology, Bangladesh Specialized Hospital, Dhaka, Bangladesh

⁶Associate Consultant, Department of Transfusion Medicine, Bangladesh Specialized Hospital, Dhaka, Bangladesh, Email: haqueanisul08@gmail.com, Orcid Id: 0000-0002-0927-4126

Corresponding Author: Dr. Ishrat Jahan, Professor, Department of Pediatrics, Bangladesh Shishu Hospital and Institute, Dhaka, Bangladesh

Abstract

Background: Typhoid fever remains a significant pediatric health burden in Bangladesh, compounded by rising antimicrobial resistance (AMR). This study aimed to describe the demographic, clinical and antimicrobial susceptibility patterns of *Salmonella Typhi* isolates in pediatric patients at a tertiary care hospital in Dhaka.

Methods: This retrospective study was carried out in the Department of Pediatrics and Neonatology, Bangladesh Specialized Hospital, Dhaka, Bangladesh and included 155 culture-confirmed pediatric typhoid cases recorded between January 2019 and December 2023. Demographic, clinical and microbiological data were analyzed using SPSS v25.

Results: Most cases occurred in children aged 9-12 years (50.3%) and 5-8 years (40.0%), with a male predominance (58.1%) and urban residence (67.7%). Resistance to first-line agents was high: Amoxiclav (62.6%), chloramphenicol (73.6%) and trimethoprim-sulfamethoxazole (97%). Fluoroquinolone and pefloxacin were resistance in 96.2% and (90%) respectively. In contrast, Sensitive to ceftriaxone (95.6%), cefixime (93.4%), azithromycin (92.3%) and meropenem (100%) remained high. Multidrug resistance was noted in 59.3% of cases, with rare XDR isolates (1.3%). Median time to defervescence was 5 days, with a mean hospital stay of 7.4 ± 2.8 days.

Conclusion: The persistence of high resistance to first-line antimicrobials and significant nalidixic acid resistant *Salmonella Typhi* (NARST) including fluoroquinolone resistance among pediatric typhoid cases in Bangladesh highlights the urgent need for revision of empirical treatment guidelines and strengthened antimicrobial resistance surveillance.

Keywords: Typhoid fever, antimicrobial resistance, pediatrics, Bangladesh, ceftriaxone.

I. INTRODUCTION

Typhoid fever remains a major global public health concern, particularly in low and middle-income countries (LMICs), where the disease continues to cause significant morbidity and mortality despite advances in sanitation, vaccination and antimicrobial therapy. Globally, it is estimated that between 11 and 20 million cases occur annually, leading to approximately 150,000 deaths, with the majority of cases caused by *Salmonella enterica* serovars Typhi and Para typhi (1). Systematic global reviews have confirmed that *S. Typhi* accounts for nearly three-quarters of all enteric fever cases, underscoring its central role in the ongoing disease burden (2,3). While typhoid fever has declined in some regions due to improved infrastructure and vaccination programs, it remains entrenched in endemic zones across South Asia and Sub-Saharan Africa, perpetuating a cycle of recurrent infections and resistance-driven treatment challenges (4,5). Children and adolescents, in particular, constitute a disproportionately high-risk group, reflecting both increased exposure to unsafe water and food and the limited reach of preventive strategies (6).

The burden of typhoid fever in South Asia is especially profound, with Bangladesh consistently

reporting among the highest incidence rates worldwide. Population-based surveillance studies, including the Surveillance for Enteric Fever in Asia Project (SEAP), demonstrated that adjusted incidence rates in urban Bangladesh exceeded 200 cases per 100,000 person-years, highlighting the intense transmission pressure in densely populated areas (7). Data from SEAP II further revealed that more than half of all confirmed enteric fever cases across South Asia originated from Bangladesh, reflecting both the endemicity and the concentration of clinical research capacity in Dhaka through icddr,b and affiliated hospitals (8). Hospital-based studies in Dhaka have consistently shown that children represent the majority of hospitalized enteric fever patients, underscoring the vulnerability of younger age groups to both infection and its complications (9). These findings align with prior prospective surveillance conducted by icddr,b, which confirmed typhoid as a persistent cause of febrile illness in Bangladesh despite national improvements in cholera control and diarrheal disease prevention (10).

Parallel to the high incidence burden, the rise of antimicrobial resistance (AMR) among typhoidal *Salmonella* has emerged as a major public health crisis. Historically, first-line agents such as chloramphenicol, ampicillin and trimethoprim-sulfamethoxazole were effective for decades; however, their efficacy has been undermined by widespread misuse and over-the-counter availability, leading to the emergence of multidrug-resistant (MDR) and XDR strains across South Asia (11). More recently, reduced susceptibility to fluoroquinolones, once the mainstay of empirical therapy, has become widespread, rendering these agents unreliable for clinical management (8). Alarming reports from Bangladesh have documented increasing azithromycin resistance, raising concerns about the potential loss of one of the last remaining effective oral therapies (9). Genomic surveillance has further confirmed the regional and intercontinental spread of these resistant strains, with resistant *S. Typhi* lineages moving between South Asian countries, amplifying the risk of outbreaks that are difficult to treat (12). Such resistance trends underscore the fragility of the current treatment arsenal, where ceftriaxone and azithromycin remain as key therapeutic options, but both face the growing threat of declining susceptibility (13).

The clinical and public health implications of these evolving resistance patterns are profound. In Bangladesh, delayed initiation of effective therapy due to empirical treatment failures has been associated with prolonged hospital stays, increased healthcare costs and worse patient outcomes (14). A cost-of-illness study within the SEAP project found that treatment of typhoid cases placed a substantial economic burden on both households and health systems, particularly when resistant infections required prolonged or alternative therapies (15). Globally, systematic reviews have shown that patients infected with resistant *S. Typhi* strains have significantly higher risks of treatment failure, complications and mortality, compared to those with susceptible infections (11). The economic consequences are equally concerning, with resistant infections linked to longer hospitalizations and increased antibiotic expenditures, straining already overburdened healthcare infrastructures in LMICs (16). The clinical challenge is exacerbated by the limited availability of new antimicrobials for enteric fever, meaning that treatment strategies rely heavily on preserving the efficacy of existing drugs.

In the present study continuous local surveillance of antibiotic sensitivity patterns is essential to guide empirical therapy, inform treatment guidelines and shape public health policies (5). Despite several multicenter studies, there remains a critical need for hospital-based research from Bangladesh to provide real-time data on resistance trends in clinical isolates (9). Such evidence is vital not only for clinical decision-making but also for national antibiotic stewardship initiatives and global strategies aimed at controlling the spread of resistant *S. Typhi*. Therefore, this study aims to evaluate the antibiotic sensitivity patterns of *Salmonella* isolates obtained from patients diagnosed with typhoid fever at a tertiary care hospital in Bangladesh, thereby contributing updated, locally relevant data to strengthen evidence-based management of this persistent and evolving public health challenge.

II. METHODS

This study employed a retrospective observational design and was conducted at Bangladesh Specialized Hospital, Dhaka, Bangladesh, over five-year period, from January 2019 to December 2023. The primary objective was to analyze the antibiotic sensitivity patterns of *Salmonella Typhi* (*S. Typhi*) isolated from patients diagnosed with typhoid fever at the hospital.

Hospital microbiology laboratory records and patient medical charts yielded 155 culture-confirmed *S. Typhi* cases, which constituted an adequate sample for statistical analysis. Only records with complete demographic, clinical and microbiological data were included in the study. Cases with incomplete records, those diagnosed as ectopic fever and those identified as *Salmonella* Para typhi A or other non-*Typhi* *Salmonella* isolates were excluded from the analysis to maintain focus on *S. Typhi* resistance trends.

Laboratory confirmation of *S. Typhi* was performed according to standard microbiological techniques, including blood culture and biochemical identification using automated and manual methods available in the hospital's microbiology department. Antibiotic susceptibility testing (AST) was conducted using the Kirby–Bauer disk diffusion method following Clinical and Laboratory Standards Institute (CLSI) guidelines.

Antibiotics tested included first-line agents (chloramphenicol, amoxiclav, trimethoprim-sulfamethoxazole), fluoroquinolones (ciprofloxacin, ofloxacin), third- generation cephalosporins (ceftriaxone, cefixime) and azithromycin, as per institutional protocols. The zone diameters were interpreted in accordance with CLSI performance standards current at the time of each test to ensure accuracy and comparability of results over the years.

All collected data were anonymized to protect patient confidentiality before analysis. Statistical analyses were performed using SPSS version 25.0 (IBM Corp., Armonk, NY, USA). Descriptive statistics were used to summarize demographic data and frequency distributions of antibiotic susceptibility. Categorical variables were expressed as frequencies and percentages. Temporal trends in antibiotic resistance patterns across the years were evaluated to observe any significant changes in sensitivity profiles. Where appropriate, chi-square tests were used to assess associations between categorical variables, with a p-value < 0.05 considered statistically significant. Socioeconomic status was classified based on monthly household income, in accordance with locally accepted economic thresholds. Participants were categorized into low, middle, and upper-middle socioeconomic groups.

Ethical approval for the study was obtained from the Institutional Review Board (IRB) of Bangladesh Specialized Hospital, with a waiver of informed consent due to the retrospective nature of the study and the use of anonymized secondary data. The study was conducted in compliance with the principles of the Declaration of Helsinki for medical research involving human subjects.

III. RESULTS

Table 1. Demographic and Baseline Characteristics of Pediatric Typhoid Cases (n = 155)

Variables	n	%
Age		
<5 year	15	9.68%
5-8 years	62	40.00%
9-12 years	78	50.32%
Sex		
Male	90	58.06%
Female	65	41.94%
Residence		
Urban	105	67.74%
Rural	50	32.26%
Socioeconomic Status		
Low Economic Class	25	16.13%
Middle Class	120	77.42%
Upper Middle Class	10	6.45%
Prior antibiotic use (within 72h)		
Yes	45	29.03%
No	110	70.97%

Among the 155 pediatric patients with culture-confirmed *Salmonella* Typhi infection, the largest age group was 9-12 years (n = 78, 50.3%), followed by 5–8 years (n = 62, 40.0%) and children younger than 5 years (n = 15, 9.7%). Males comprised a slight majority of cases (n = 90, 58.1%) compared with females (n = 65, 41.9%). Most patients resided in urban areas (n = 105, 67.7%), while 32.3% (n = 50) were from rural regions. In terms of socioeconomic distribution, the middle-class group predominated (n = 120, 77.4%), followed by the low economic class (n = 25, 16.1%) and a smaller proportion from the upper-middle class (n = 10, 6.5%). Regarding recent medication history, 29.0% (n = 45) of patients had a history of antibiotic use within 72 hours prior to hospital presentation, while the majority (71.0%, n = 110) had no prior antibiotic exposure.

Table 2. Clinical Presentation and Complications of Pediatric Typhoid Cases (n = 155)

Variables	n	%
Fever duration before admission		
≤7 days	43	27.74%
8–14 days	65	41.94%
>14 days	47	30.32%
Other clinical features		
Abdominal pain	126	81.29%
Diarrhea	72	46.45%
Vomiting	64	41.29%
Hepatomegaly	52	33.55%

Among the 155 pediatric patients, the duration of fever before hospital admission was 8-14 days in the largest proportion (n = 65, 41.9%), followed by more than 14 days in 47 cases (30.3%) and 7 days or less in 43 cases (27.7%). The most frequent clinical feature was abdominal pain, reported in 126 patients (81.3%), followed by diarrhea in 72 patients (46.5%) and vomiting in 64 patients (41.3%). Hepatomegaly was documented in 52 children (33.6%).




Figure 1: Distribution of Salmonella Isolates (n = 155)

The yearly distribution of culture-confirmed Salmonella Typhi isolates demonstrated relatively consistent numbers across the five years of surveillance. In 2019, there were 24 isolates (15.2%), increasing slightly to 29 (19.0%) in 2020 and 34 (21.9%) in 2021. The highest number was observed in 2022, with 35 isolates (22.9%), followed by 33 cases (21.3%) in 2023. Overall, the data indicate a steady burden of confirmed pediatric typhoid cases over the five-year period.

Table 3. Antibiotic Susceptibility Profile of S. Typhi Isolates (n = 155)

Antibiotic	Sensitive n (%)	Intermediate n (%)	Resistant n (%)
Amoxiclav	20 (37.4)	-	135 (62.6)
Chloramphenicol	41 (26.4)	-	114 (73.6)
TMP-SMX	5 (3)	-	150 (97)
Ciprofloxacin	16 (10)	-	139 (90)
Ceftriaxone	148 (95.6)	-	7 (4.4)

Cefixime	145 (93.4)	–	10 (6.6)
Azithromycin	143 (92.3)	–	12(7.7)
Nalidixic Acid	06(3.8)		149(96.2)
Pefloxacin	16 (10)	-	139(90)
Meropenem	155 (100.0)	–	0 (0.0)

Among the 155 *Salmonella Typhi* isolates, meropenem demonstrated 100% susceptibility (n = 155, 100.0%) with no intermediate or resistant isolates detected. High susceptibility was also observed for ceftriaxone (n = 148, 95.6%) and cefixime (n = 145, 93.4%), with resistance noted in only 7 (4.4%) and 10 (6.6%) isolates, respectively. Azithromycin retained high activity, with 143 isolates (92.3%) sensitive and 12 isolates (7.7%) resistant. In contrast, marked resistance was observed to first-line antibiotics. Amoxiclav showed sensitivity in only 20 isolates (37.4%), with resistance in 135(62.6%) isolates. Chloramphenicol susceptibility was 26.4% (n = 41), with 73.6% (n = 114) resistant, and trimethoprim-sulfamethoxazole (TMP-SMX) was sensitive in 3% (n = 5) and resistant in 97.9% (n = 149). Fluoroquinolone (Ciprofloxacin, Pefloxacin) exhibited 10% (n = 16) sensitive and 90% (n = 139) resistant isolates. Nalidixic acid resistant *Salmonella Typhi* (NARST) isolates exhibited resistance to fluoroquinolones in vivo.




Figure 2: Multidrug-Resistance (MDR) and Extensively-Drug Resistance (XDR) Patterns (n = 155)

Among the 155 *Salmonella Typhi* isolates, multidrug resistance (MDR) defined as concurrent resistance to ampicillin, chloramphenicol, and trimethoprim-sulfamethoxazole (TMP-SMX) was observed in 45 isolates (29.0%). Fluoroquinolone (FQ) resistance was the most common resistance, present in 85 isolates (54.8%). Extensively drug-resistant (XDR) strains, characterized by resistance to first-line agents, fluoroquinolones, and third-generation cephalosporins, were identified in 2 isolates (1.3%), indicating that while rare, XDR *S. Typhi* is present in this cohort.




Figure 3: Treatment Patterns and Clinical Outcomes of Pediatric Typhoid Cases (n = 155)

Among the 155 pediatric typhoid cases, the most common empiric therapy administered was ceftriaxone, used in 100 patients (64.5%), followed by azithromycin in 29 patients (18.7%), combination regimens in 12 patients (7.7%) and other antibiotic choices in 14 patients (9.0%). For definitive therapy, ceftriaxone remained the predominant agent (61.3%, n = 95), while azithromycin was used in 40 patients (25.8%). Therapy adjustments based on culture and sensitivity results occurred in 13 patients (8.4%), whereas the majority (65.8%, n = 102) continued their initial empiric regimen. The median time to defervescence following initiation of treatment was 5 days (interquartile range [IQR]: 3–7 days) and the mean length of hospital stay was 7.4 ± 2.8 days.

IV. DISCUSSION

This retrospective analysis of 155 pediatric cases of culture-confirmed *Salmonella Typhi* (S. Typhi) infection in a tertiary care hospital in Dhaka, Bangladesh, offers a comprehensive overview of demographic distribution, clinical profiles, antimicrobial susceptibility patterns, resistance phenotypes, treatment modalities and outcomes over a five-year period. The findings underscore critical public health concerns around antimicrobial resistance (AMR) in pediatric typhoid fever, while highlighting the efficacy of current treatment protocols and the importance of continuous surveillance.

The demographic profile of affected children revealed a predominance in the 9-12-year age group (50.3%), followed by the 5-8-year cohort (40%). A male preponderance (58.1%) and a substantial urban concentration (67.7%) were also noted. These trends are consistent with regional patterns reported in Dhaka and other South Asian urban centers. For instance, a multicenter study by Qamar et al. under the SEAP project similarly identified higher incidence rates in school-aged children in urban settings, aligning with the age and residential trends observed in our cohort (8). Prior antibiotic use within 72 hours of admission was documented in 29%, a figure supported by Hamdulay et al., who highlighted the widespread unsupervised antibiotic usage as a contributor to AMR in LMICs (17).

Clinically, the majority of children presented after more than one week of febrile illness, with 41.9% presenting between 8-14 days and 30.3% after 14 days. Abdominal pain (81.3%), diarrhea (46.5%) and vomiting (41.3%) were among the most frequent symptoms, while hepatomegaly was noted in one-third of cases. This symptom distribution and the low rate of complications are closely aligned with findings from Nepal and India, where similar symptom prevalence and favorable outcomes have been reported in pediatric patients (18,19).

The annual distribution of isolates remained relatively consistent across the five-year study period, suggesting a sustained endemic burden. This steady trend reflects regional surveillance reports, such as the SEAP study, which also observed persistent incidence rates in Bangladesh over similar timeframes (8).

A major highlight of the current study is the concerning level of resistance to first-line antibiotics. Resistance to amoxiclav (62.6%), chloramphenicol (73.6%) and TMP-SMX (97%) was observed, with fluoroquinolone resistance accounting for a combined 90% of isolates. These figures mirror national and regional studies. Tanmoy et al., in a 24- year resistance trend analysis, similarly reported resistance rates exceeding 59.3% for first-line agents (14). Additionally, SEAP data indicated fluoroquinolone non-susceptibility as a growing concern in South Asia (8). These findings collectively underscore the declining reliability of older antibiotics and the urgent need for treatment reassessment.

In contrast, third-generation cephalosporins and azithromycin demonstrated high efficacy in our study: ceftriaxone (95.6%), cefixime (93.4%), azithromycin (92.3%) and meropenem (100%). These findings are supported by multiple clinical and surveillance studies. For instance, the Cochrane review by Kuehn et al. reaffirmed ceftriaxone's role as a cornerstone of typhoid therapy, while Hye et al. observed similarly high ceftriaxone efficacy in pediatric patients in Dhaka (20,21). The preserved sensitivity to these agents justifies their continued use as empiric and definitive therapies, as supported by global pediatric treatment guidelines (19).

Multidrug-resistant (MDR) *S. Typhi* strains, defined as resistance to all three first-line antibiotics, were found in 59.3% of cases in our cohort. NARST associated fluoroquinolone resistance was higher at 90%, while extensively drug-resistant (XDR) isolates resistant to first-line agents, fluoroquinolones and third-generation cephalosporins were rare (1.3%). These figures are closely aligned with those reported by Ghurnee et al. and Tanmoy et al., who noted MDR prevalence between 25–35% and sporadic emergence of XDR strains in Bangladesh (22,23). Although infrequent, the detection of XDR isolates warrants proactive surveillance and restriction of empiric use of cephalosporins without culture confirmation.

In terms of treatment practices, ceftriaxone was the predominant empiric (64.5%) and definitive (61.3%) therapy. Azithromycin was used in 18.7% empirically and 25.8% definitively. The median time to defervescence was five days (IQR 3–7) and the mean hospital stay was 7.4 ± 2.8 days. These figures compare well with those in clinical trials from South Asia, where median defervescence ranged from 4–6 days and length of stay hovered around 6–8 days (18,24). Adjustment of treatment post-culture was necessary in 8.4% of cases, indicating the importance of microbiological guidance in managing resistant strains. Favorable clinical outcomes were observed overall, with as reported by Memon et al. and Parry et al (13,19).

In summary, this study reaffirms the high burden of pediatric typhoid fever in Bangladesh, coupled with alarming levels of resistance to conventional antibiotics. While newer agents like ceftriaxone, azithromycin and meropenem remain effective, the detection of MDR and XDR strains highlights the pressing need for stringent antimicrobial stewardship, robust surveillance systems and community-level education to curb inappropriate antibiotic use and mitigate the risk of further resistance evolution.

Limitations of The Study

The study was conducted in a single hospital with a small sample size. So, the results may not represent the whole community.

V. CONCLUSION

This retrospective study highlights the persistent burden of pediatric typhoid fever in Bangladesh, characterized by a predominance in school-aged children and a significant urban representation. High levels of resistance to first-line agents' amoxiclav, chloramphenicol and trimethoprim-sulfamethoxazole alongside widespread quinolone and NARST associated fluoroquinolone resistant, underline the diminishing utility of these antibiotics in routine clinical practice. The preserved susceptibility to ceftriaxone, cefixime, azithromycin and meropenem supports their continued use as empiric and definitive therapies in clinically suspected typhoid fever cases. The detection of rare but emerging extensively drug-resistant (*S. Typhi*) isolates emphasizes the need for robust surveillance, prudent antibiotic stewardship and region-specific treatment guidelines to prevent further escalation of resistance. Favorable clinical outcomes, with rapid defervescence, zero relapse and mortality, reflect effective therapeutic strategies but also

highlight the importance of ongoing monitoring and adaptation of treatment protocols in response to evolving resistance trends.

Funding: No funding sources

Conflict of interest: None declared

Ethical approval: The study was approved by the Institutional Ethics Committee

REFERENCES

- [1]. Stanaway JD, Reiner RC, Blacker BF, Goldberg EM, Khalil IA, Troeger CE, et al. The global burden of typhoid and paratyphoid fevers: a systematic analysis for the Global Burden of Disease Study 2017. *The Lancet Infectious Diseases*. 2019 Apr 1;19(4):369–81.
- [2]. Buckle GC, Walker CLF, Black RE. Typhoid fever and paratyphoid fever: Systematic review to estimate global morbidity and mortality for 2010. *J Glob Health*. 2012 June;2(1):010401.
- [3]. Als D, Radhakrishnan A, Arora P, Gaffey MF, Campisi S, Velummailum R, et al. Global Trends in Typhoidal Salmonellosis: A Systematic Review. *Am J Trop Med Hyg*. 2018 Sept;99(3_Suppl):10–9.
- [4]. Marchello CS, Hong CY, Crump JA. Global Typhoid Fever Incidence: A Systematic Review and Meta-analysis. *Clin Infect Dis*. 2019 Mar 7;68(Suppl 2):S105–16.
- [5]. Mogasale V, Maskery B, Ochial RL, Lee JS, Mogasale VV, Ramani E, et al. Burden of typhoid fever in low-income and middle-income countries: a systematic, literature-based update with risk-factor adjustment. *Lancet Glob Health*. 2014 Oct;2(10):e570–580.
- [6]. Wang H, Zhang P, Zhao Q, Ma W. Global burden, trends and inequalities for typhoid and paratyphoid fever among children younger than 15 years over the past 30 years. *J Travel Med*. 2024 Dec 10;31(8):taae140.
- [7]. Garrett DO, Longley AT, Aiemjoy K, Yousafzai MT, Hemlock C, Yu AT, et al. Incidence of typhoid and paratyphoid fever in Bangladesh, Nepal, and Pakistan: results of the Surveillance for Enteric Fever in Asia Project. *Lancet Glob Health*. 2022 July;10(7):e978–88.
- [8]. Qamar FN, Yousafzai MT, Dehraj IF, Shakoor S, Irfan S, Hotwani A, et al. Antimicrobial Resistance in Typhoidal Salmonella: Surveillance for Enteric Fever in Asia Project, 2016–2019. *Clin Infect Dis*. 2020 Dec 1;71(Suppl 3):S276–84.
- [9]. Saha S, Sayeed KMI, Saha S, Islam MS, Rahaman A, Islam M, et al. Hospitalization of Pediatric Enteric Fever Cases, Dhaka, Bangladesh, 2017–2019: Incidence and Risk Factors. *Clin Infect Dis*. 2020 Dec 1;71(Suppl 3):S196–204.
- [10]. Darton TC, Meiring JE, Tonks S, Khan MA, Khanam F, Shakya M, et al. The STRATAA study protocol: a programme to assess the burden of enteric fever in Bangladesh, Malawi and Nepal using prospective population census, passive surveillance, serological studies and healthcare utilisation surveys. *BMJ Open*. 2017 July 2;7(6):e016283.
- [11]. Browne AJ, Kashef Hamadani BH, Kumaran EAP, Rao P, Longbottom J, Harriss E, et al. Drug-resistant enteric fever worldwide, 1990 to 2018: a systematic review and meta-analysis. *BMC Med*. 2020 Jan 3;18(1):1.
- [12]. da Silva KE, Tanmoy AM, Pragasam AK, Iqbal J, Sajib MSI, Mutreja A, et al. The international and intercontinental spread and expansion of antimicrobial-resistant *Salmonella* Typhi: a genomic epidemiology study. *Lancet Microbe*. 2022 Aug;3(8):e567–77.
- [13]. Memon H, Saeed F, Iqbal M, Saboohi E, Hanif S, Mallick AHH. Association of extensively drug resistant salmonella infection in children with typhoid fever. *Pak J Med Sci*. 2022;38(7):1864–9.
- [14]. Tanmoy AM, Hooda Y, Sajib MSI, Rahman H, Sarkar A, Das D, et al. Trends in antimicrobial resistance amongst *Salmonella* Typhi in Bangladesh: A 24-year retrospective observational study (1999–2022). *PLOS Neglected Tropical Diseases*. 2024 Oct 4;18(10):e0012558.
- [15]. Mejia N, Pallas SW, Saha S, Udin J, Sayeed KMI, Garrett DO, et al. Typhoid and Paratyphoid Cost of Illness in Bangladesh: Patient and Health Facility Costs From the Surveillance for Enteric Fever in Asia Project II. *Clin Infect Dis*. 2020 Dec 1;71(Suppl 3):S293–305.
- [16]. Kim C, Frost I, Naylor NR, Au H, Kim Y, Lee Y, et al. Length of hospital stay and associated treatment costs for patients with susceptible and antibiotic-resistant *Salmonella* infections: a systematic review and meta-analysis. *BMJ Open*. 2025 June 23;15(6):e092494.
- [17]. Hamdulay K, Rawekar R, Tayade A, Kumar S, Acharya S. Evolving Epidemiology and Antibiotic Resistance in Enteric Fever: A Comprehensive Review. *Cureus*. 16(6):e63070.
- [18]. Basnyat B, Giri A, Karkey A, Dongol S, Arjyal A, Maharjan A, et al. Azithromycin and cefixime combination versus azithromycin alone for the out-patient treatment of clinically suspected or confirmed uncomplicated typhoid fever in South Asia: A randomised controlled trial protocol. *Wellcome Open Research*. 2021;6:207.
- [19]. Parry CM, Qamar FN, Rijal S, McCann N, Baker S, Basnyat B. What Should We Be Recommending for the Treatment of Enteric Fever? *Open Forum Infect Dis*. 2023 May;10(Suppl 1):S26–31.
- [20]. Kuehn R, Stoesser N, Eyre D, Darton TC, Basnyat B, Parry CM. Treatment of enteric fever (typhoid and paratyphoid fever) with cephalosporins. *Cochrane Database Syst Rev*. 2022 Nov 24;2022(11):CD010452.
- [21]. Hye MA, Afroza S, Kabir AL, Habib RB, Khanom ME. Comparing the Efficacy of Amoxicillin and Ceftriaxone in Clinical Management of Uncomplicated Enteric Fever in Children: A Cost-effective Approach in Bangladesh. *The Journal of Ad-din Women's Medical College*. 2023 Jan 8;11(1):14–21.
- [22]. Ghurnee O, Ghosh AK, Abony M, Aurin SA, Fatema AN, Banik A, et al. Isolation of Multi-Drug Resistant (MDR) and Extensively Drug Resistant (XDR) *Salmonella* typhi from Blood Samples of Patients Attending Tertiary Medical Centre in Dhaka City, Bangladesh. *Advances in Microbiology*. 2021 Sept 2;11(9):488–98.
- [23]. Tanmoy AM, Westeel E, De Bruyne K, Goris J, Rajoharison A, Sajib MSI, et al. *Salmonella enterica* Serovar Typhi in Bangladesh: Exploration of Genomic Diversity and Antimicrobial Resistance. *mBio*. 2018 Nov 13;9(6):e02112–18.
- [24]. Siddiqi FG, Rais H, Zafar F, Bina, Anwar T, Ally Z, et al. Treatment of 21st-Century typhoid fever in children; open-label mono versus combination drug therapy: A multicenter randomized control study. *International Journal of Endorsing Health Science Research*. 2021 Sept 1;9(3):384–94.