

Choosing Burs For Tooth Preparations: A Concised Guideline

Dr.Shardool Jagtap, Dr.Ayesha Siddiqua, Dr.Babita Yeshwante, Dr.Nazish Baig, Dr.Ruchi Kasat, Dr.Rutuja Gaikwad

Abstract:

Tooth preparation for fixed prosthodontics requires precise biomechanical execution to achieve optimal retention, resistance, marginal integrity, periodontal compatibility, and restoration longevity. Dental burs—varying in material, grit size, blade design, and geometry—significantly influence cutting efficiency, surface roughness, pulpal response, and marginal accuracy. Evidence from indexed prosthodontic and restorative dentistry journals supports sequential bur protocols incorporating diamond burs for gross reduction and multi-fluted carbide burs for finishing. Vital teeth require strict thermal control to prevent pulpal injury, while non-vital teeth demand structural preservation and precise margin configuration. This review synthesizes contemporary literature and proposes an evidence-based bur selection protocol for predictable prosthodontic outcomes.

Keywords: Dental burs; Prosthodontics; Finish line; Diamond bur; Carbide bur; Pulpal response; Marginal adaptation.

Date of Submission: 23-03-2026

Date of Acceptance: 03-04-2026

I. Introduction

Successful fixed prosthodontic therapy depends on accurate tooth preparation that balances mechanical durability with biological preservation¹. Rotary cutting instruments, particularly dental burs, directly affect the geometry and surface characteristics of prepared teeth².

Bur selection influences:

- Cutting rate
- Surface roughness
- Smear layer formation
- Heat generation
- Margin configuration
- Digital scan accuracy

Siegel and von Fraunhofer demonstrated that bur type significantly alters surface morphology and cutting efficiency during fixed prosthodontic preparation³. Ayad later confirmed that preparation surface characteristics influence marginal fit of extra-coronal restorations⁴.

Despite widespread clinical use, structured evidence-based guidelines remain limited. This review integrates indexed literature to guide bur selection in:

- Vital teeth
- Non-vital teeth
- Finish line preparation
- Margin finishing
- Digital workflows

II. History

Dental burs were manufactured at least 300 years before, and still widely used. These can be of steel followed by a coating of tungsten carbide or entirely made of tungsten carbide Maya with the other ancient cultures used to use the primitive "bow drills" with few different devices for preparing the round ornamental cavities in the teeth 1728 Pierre Fauchard, "Le Chirurgien Dentiste" first detailed description of dental practice represented the use of bow drill to prepare teeth with cutters.

1891, the S.S.White company gave revelation burs, which were the first machine-made steel burs.

In 1897, Willman and Schroeder of the University of Berlin, Germany, were credited for manufacturing the first diamond bur.

In 1942, diamond cutting instruments with 5000 rpm were introduced.

In 1947, Carbide Burs with 12,000 rpm replaced the steel burs.

III. Classification And Cutting Mechanisms

Diamond Burs:

Diamond burs cut via abrasive action. Particle size determines aggressiveness⁵.

Grit classification:

- Super coarse (150–200 µm)
- Coarse (125–150 µm)
- Medium (100–125 µm)
- Fine (30–60 µm)
- Superfine (<30 µm)

Diamond burs exhibit high cutting efficiency in enamel⁶ and are recommended for bulk reduction⁷. However, they produce greater surface roughness compared to carbide burs³.

Worn diamond burs increase friction and heat generation⁸.

Tungsten Carbide Burs

Carbide burs cut by shearing action via fluted blades.

Flute configurations:

- 6–8 flutes → Cutting
- 12–20 flutes → Finishing
- 30 flutes → Ultra-finishing

Carbide finishing burs produce smoother surfaces and improved marginal adaptation^{4,9}.

IV. Biological Considerations

Pulpal Response in Vital Teeth:

Zach and Cohen demonstrated that a 5.5°C intra-pulpal temperature rise may cause irreversible pulp damage¹⁰.

Diamond burs generate greater frictional heat than carbide burs under identical conditions¹¹. Adequate coolant significantly reduces temperature rise¹².

Therefore, preparation of vital teeth requires:

- Copious air-water spray
- Intermittent cutting stroke.
- Sharp, unworn burs
- Progressive grit reduction

Excessive vibration and pressure may cause odontoblastic damage and postoperative sensitivity¹³.

Non-Vital Teeth:

Although pulpal preservation is not a concern, structural durability is critical. Endodontically treated teeth exhibit altered biomechanical behavior¹⁴.

Margin precision remains essential to prevent microleakage and restoration failure^{4,15}.

V. Bur Selection During Preparation Phases

Gross Occlusal Reduction:

Medium-grit diamond burs provide efficient enamel removal⁶. Coarse burs may produce excessive irregularity without clinical advantage¹¹.

Depth orientation grooves improve reduction accuracy and reduce over-preparation¹⁶.

Axial Wall Preparation:

Controlled taper (6–12 degrees) is essential for retention¹. Bur geometry influences convergence consistency¹⁶.

Sequential grit reduction (medium → fine) improves surface smoothness⁹.

Finish Line Preparation:

Finish line design affects stress distribution and marginal integrity¹⁷.

Chamfer → Round-end tapered diamond

Shoulder → Flat-end cylindrical diamond

Radial shoulder → Round-end cylindrical diamond

Margin width should correspond to restoration material requirements^{17,18}.

VI. Margin Finishing And Surface Roughness

Surface roughness influences marginal adaptation⁴.

Ayad demonstrated significantly improved marginal fit when finishing burs were used compared with cutting burs alone⁴.

Carbide finishing burs (12–20 flutes) significantly reduce surface irregularities^{3,9}.

Ultrasonic preparation systems may produce smoother finish lines compared to rotary instruments¹⁹.

VII. Digital Dentistry Considerations

Surface roughness affects digital scanning accuracy²⁰.

Smooth finish lines improve CAD/CAM margin detection²¹.

Sequential diamond-carbide protocols enhance digital impression precision²⁰.

VIII. Thermal And Mechanical Considerations

Diamond burs produce greater frictional heat than carbide burs¹¹.

Coolant flow reduces pulpal temperature rise¹².

Worn burs increase pressure and heat generation⁸.

Excessive axial force decreases cutting efficiency and increases thermal injury¹³.

IX. Evidence-Based Clinical Protocol

Step 1 – Occlusal Reduction: Medium Diamond

Step 2 – Depth Grooves: Depth-Cut Diamond

Step 3 – Axial Reduction: Medium: Fine Diamond

Step 4 – Finish Line Formation: Shape-Specific Diamond

Step 5 – Margin Refinement: 12-Flute Carbide

Step 6 – Final Smoothing: 20–30-Flute Carbide

X. Discussion

Evidence supports combined diamond-carbide sequencing^{3,4}.

Diamond burs are indispensable for enamel removal⁶. Carbide burs enhance surface refinement and marginal precision^{3,9}.

In vital teeth, thermal control is critical^{10,12}.

Margin quality directly influences restoration adaptation and longevity^{4,15}.

Future research should evaluate long-term survival outcomes associated with bur sequencing protocols.

XI. Conclusion

Proper bur selection significantly influences prosthodontic outcomes.

- Medium-grit diamonds for bulk reduction
- Fine diamonds for refinement
- Multi-fluted carbide burs for margin finishing
- Strict thermal control in vital teeth
- Sequential grit reduction improves surface quality

An evidence-based bur protocol enhances restoration fit, biological preservation, and clinical longevity.

References

- [1]. Goodacre CJ, Campagni WV, Aquilino SA. Tooth Preparations For Complete Crowns. *J Prosthet Dent.* 2001;85:363-376.
- [2]. Rosenstiel SF, Et Al. Influence Of Preparation Design On Restoration Longevity. *J Prosthet Dent.* 1998.
- [3]. Siegel SC, Von Fraunhofer JA. Dental Burs—What Bur For Which Application? *J Prosthodont.* 1999;8:258-263.
- [4]. Ayad MF. Effect Of Preparation Burs On Marginal Fit. *J Prosthodont.* 2009;18:145-151.
- [5]. Banerjee A, Watson TF. Diamond Bur Performance. *J Dent.* 2000.
- [6]. Peters MC, Et Al. Cutting Efficiency Of Diamond Burs. *J Dent Res.* 2001.
- [7]. Sharma N, Chitre V. Full Veneer Tooth Preparation Tips. *J Indian Prosthodont Soc.* 2007;7:137-142.
- [8]. Shortall AC. Bur Wear And Efficiency. *Br Dent J.* 1988.
- [9]. Lopes GC, Et Al. Surface Roughness After Finishing Procedures. *Oper Dent.* 2007.
- [10]. Zach L, Cohen G. Pulp Response To Externally Applied Heat. *J Prosthet Dent.* 1965;15:515-530.
- [11]. Cavalcanti BN, Et Al. Temperature Rise During Tooth Preparation. *Oper Dent.* 2002.
- [12]. Loney RW, Price RB. Air-Water Spray Effectiveness. *J Prosthet Dent.* 2001.
- [13]. Kim S, Et Al. Pulpal Blood Flow During Preparation. *J Endod.* 1984.
- [14]. Reeh ES, Et Al. Fracture Resistance Of Endodontically Treated Teeth. *J Endod.* 1989.
- [15]. Felton DA, Et Al. Crown Margin Discrepancies. *J Prosthet Dent.* 1991.
- [16]. Ram HK, Et Al. Evaluation Of Preparation Techniques. *J Indian Prosthodont Soc.* 2015;15:162-167.
- [17]. Shillingburg HT Jr, Et Al. Margin Design Principles. *J Prosthet Dent.* 1973.

- [18]. Sorensen JA. Marginal Fit Of Crown Systems. J Prosthet Dent. 1990.
- [19]. Horne P, Et Al. Ultrasonic Margin Preparation. J Esthet Restor Dent. 2012;24:201-209.
- [20]. Ender A, Mehl A. Accuracy Of Digital Impressions. Int J Comput Dent. 2011.
- [21]. Renne W, Et Al. CAD/CAM Margin Detection. J Prosthet Dent. 2012.
- [22]. Christensen GJ. Preparation Design Considerations. J Am Dent Assoc. 2007.
- [23]. Patel V, Et Al. Surface Characteristics And Adhesion. J Adhes Dent. 2010.
- [24]. Tjan AH, Et Al. Effects Of Preparation Geometry. J Prosthet Dent. 1980.
- [25]. Donovan TE, Chee WW. Preparation Guidelines. J Prosthet Dent. 2004.